

BUILDING BLOCKS OF ÉTALE ENDOMORPHISMS OF COMPLEX PROJECTIVE MANIFOLDS

NOBORU NAKAYAMA AND DE-QI ZHANG

ABSTRACT. Étale endomorphisms of complex projective manifolds are constructed from two building blocks up to isomorphism if the good minimal model conjecture is true. They are the endomorphisms of abelian varieties and the nearly étale rational endomorphisms of weak Calabi-Yau varieties.

1. INTRODUCTION

We work over the field $\mathbb{C}$ of complex numbers. In this paper, we shall give a systematic study of étale endomorphisms of nonsingular projective varieties. The étaleness assumption is quite natural because every surjective endomorphism of X is étale provided that X is a nonsingular projective variety and is non-uniruled (cf. Remark 1.1 below). In the study of birational classification of algebraic varieties, we usually have the following three reductions, where κ denotes the Kodaira dimension and q the irregularity:

- (A) Varieties of $\kappa > 0 \Rightarrow$ varieties of $\kappa = 0$, by the Iitaka fibration.
- (B) Varieties of $\kappa = 0 \Rightarrow$ abelian varieties and varieties with $\kappa = q = 0$, by the Albanese map.
- (C) Uniruled varieties $\Rightarrow$ non-uniruled varieties, by the maximal rationally connected fibration (cf. [10] and [40]).

We want to show that there are similar reductions in the study of étale endomorphisms of nonsingular projective varieties. Theorems A, B, and C below correspond to the reductions (A), (B), and (C), respectively. See [57] for the case of automorphisms.

1.1. The reduction (A). This reduction is based on the Iitaka fibration. Let X be a nonsingular projective variety with $\kappa(X) > 0$ and $f: X \rightarrow X$ a surjective morphism. From a standard argument of pluricanonical systems, we infer that f induces a birational automorphism g of the base space Y of the Iitaka fibration $X \cdots \rightarrow Y$. Theorem A below shows that the order of g is finite. This was conjectured in several papers (cf.

2000 *Mathematics Subject Classification.* 14E20, 14E07, 32H50.

Key words and phrases. endomorphism, Iitaka fibration, Albanese map, rationally connected variety.

The first named author is partly supported by the Grant-in-Aid for Scientific Research (C), Japan Society for the Promotion of Science. The second named author is supported by an ARF of NUS.

[1, Proposition 6.4], [21, Proposition 3.7]). Theorem A treats not only holomorphic surjective endomorphisms of projective varieties of $\kappa > 0$ but also dominant meromorphic endomorphisms of compact complex manifolds of $\kappa > 0$ in the class $\mathcal{C}$ in the sense of Fujiki [18]. Note that a compact complex manifold is in the class $\mathcal{C}$ if and only if it is bimeromorphic to a compact Kähler manifold (cf. [53]).

Theorem A. *Let X be a compact complex manifold in the class $\mathcal{C}$ of $\kappa(X) \geq 1$ and let $f: X \dashrightarrow X$ be a dominant meromorphic map. Let W_m be the image of the m -th pluricanonical map*

$$\Phi_m: X \dashrightarrow |mK_X|^\vee = \mathbb{P}(H^0(X, mK_X))$$

for a positive integer m with $|mK_X| \neq \emptyset$. Then there is an automorphism g of W_m of finite order such that $\Phi_m \circ f = g \circ \Phi_m$.

Remark. (1) If f is holomorphic, then, resolving the indeterminacy locus of Φ_m , we may assume that both $f: X \rightarrow X$ and $\Phi_m: X \rightarrow W_m$ are holomorphic so that $\Phi_m \circ f = g \circ \Phi_m$. This is because f is étale and we can take an equivariant resolution (in the sense of Section 1.4 below) of the graph of Iitaka fibration (cf. the proof of Lemma 5.2).

(2) Theorem A is known to be true by Deligne and Nakamura-Ueno when X is Moishezon and f is a bimeromorphic automorphism (cf. [52, Theorem 14.10], [47]).

1.2. The reduction (B). For a compact Kähler manifold M with $c_1(M)_{\mathbb{R}} = 0$, we have a finite étale cover $\widetilde{M} \rightarrow M$ such that $\widetilde{M} \simeq T \times F$ for a complex torus T and a simply connected manifold F with $c_1(F) = 0$, by Bogomolov's decomposition theorem (cf. [8], [4]). For a normal projective variety V with only canonical singularities and with torsion K_V , we have the following weak decomposition by Kawamata [33, Corollary 8.4]: *There exists a finite étale covering $F \times A \rightarrow V$ for a weak Calabi-Yau variety F and an abelian variety A .* Here, a normal projective variety F is called *weak Calabi-Yau* if F has only canonical singularities, $K_F \sim_{\mathbb{Q}} 0$, and

$$q^{\max}(F) := \max\{q(F') \mid F' \rightarrow F \text{ is finite étale}\} = 0$$

(cf. Section 4.1). If F is a nonsingular weak Calabi-Yau variety, then $\pi_1(F)$ is finite by Bogomolov's decomposition theorem, so a finite étale cover of F is expressed as a product of holomorphic symplectic manifolds and Calabi-Yau manifolds.

In order to study the surjective endomorphisms of a nonsingular projective variety X with $\kappa(X) = 0$, we assume the existence of a good minimal model V of X ; but we allow the variety V to have canonical singularities as in [33]. Then it has a meaning to consider the reduction of the endomorphisms to those of weak Calabi-Yau varieties F and those

of abelian varieties A by an étale covering $F \times A \rightarrow V$. Unfortunately, a holomorphic surjective endomorphism of X induces only a rational map $V \dashrightarrow V$, but it satisfies the condition of *nearly étale map*, which is introduced in Section 3. Therefore, Theorem B below is meaningful for the reduction of type (B):

Theorem B. *Let V be a normal projective variety with only canonical singularities such that $K_V \sim_{\mathbb{Q}} 0$. Let $h: V \dashrightarrow V$ be a dominant rational map which is nearly étale. Then there exist an abelian variety A , a weak Calabi-Yau variety F , a finite étale morphism $\tau: F \times A \rightarrow V$, a nearly étale dominant rational map $\varphi_F: F \dashrightarrow F$, and a finite étale morphism $\varphi_A: A \rightarrow A$ such that $\tau \circ (\varphi_F \times \varphi_A) = h \circ \tau$, i.e., the diagram below is commutative:*

$$\begin{array}{ccc} F \times A & \xrightarrow{\varphi_F \times \varphi_A} & F \times A \\ \tau \downarrow & & \downarrow \tau \\ V & \xrightarrow{h} & V. \end{array}$$

Remark. (1) By the proof, the commutative diagram above is birationally Cartesian.

(2) If the algebraic fundamental group $\pi_1^{\text{alg}}(F)$ is finite, then φ_F is a birational automorphism (cf. Section 4.4). In particular, if V has only terminal singularities and $q^{\text{max}}(V) = \dim V - 2$, then φ_F is an automorphism, since F is a K3 surface or an Enriques surface.

1.3. The reduction (C). Let X be a uniruled nonsingular projective variety. A *maximal rationally connected fibration* of X in the sense of [10] and [40] is obtained by a certain rational map $X \dashrightarrow \text{Chow}(X)$ into the Chow variety $\text{Chow}(X)$, which assigns a general point $x \in X$ a maximal rationally connected subvariety containing x . Let Y be the normalization of the image of $X \dashrightarrow \text{Chow}(X)$ and let $\pi: X \dashrightarrow Y$ be the induced rational fibration. Assume that X admits an étale endomorphism $f: X \rightarrow X$. Then there is an endomorphism $h: Y \rightarrow Y$ such that $\pi \circ f = h \circ \pi$ (cf. Lemma 5.2). Since rationally connected manifolds are simply connected, the endomorphism f is induced from h . In Theorem C below, we shall show that h is nearly étale.

Theorem C. *Let X be a projective manifold with an étale endomorphism f . Assume that X is uniruled. Then there exist a projective manifold M with an étale endomorphism $f_M: M \rightarrow M$, a non-uniruled normal projective variety Y with a nearly étale endomorphism $h: Y \rightarrow Y$, a birational morphism $\mu: M \rightarrow X$, and a surjective morphism $\pi: M \rightarrow Y$ such that*

- (1) $\pi \circ f_M = h \circ \pi$, $\mu \circ f_M = f \circ \mu$, i.e., the diagram below is commutative:

$$\begin{array}{ccccc} Y & \xleftarrow{\pi} & M & \xrightarrow{\mu} & X \\ h \downarrow & & f_M \downarrow & & f \downarrow \\ Y & \xleftarrow{\pi} & M & \xrightarrow{\mu} & X, \end{array}$$

- (2) $\deg f = \deg f_M = \deg h$,
 (3) $\pi \circ \mu^{-1}: X \dashrightarrow M \rightarrow Y$ is birational to the maximal rationally connected fibration of X .

Remark.

- (1) $h^{-1}(Y_{\text{rat}}) = Y_{\text{rat}}$ and the restriction $Y_{\text{rat}} \rightarrow Y_{\text{rat}}$ of h is étale for the open subset $Y_{\text{rat}} \subset Y$ consisting of the smooth points and the points of rational singularity, by Proposition 3.11 and Proposition 3.10, (2).
 (2) If Y has the relative canonical model Y_{can} for resolutions of singularities of Y , then, by Lemma 3.8, h lifts to an étale endomorphism of Y_{can} and also to an étale endomorphism of a certain resolution Y' of singularities of Y . The recent paper [5] has announced a proof of the existence of minimal models of varieties of general type even in a relative setting. The existence of the relative canonical model Y_{can} follows from the result.

1.4. Equivariant resolutions. Let V be a normal projective complex variety and $f: V \rightarrow V$ an étale endomorphism. Then there exists a resolution $\mu: X \rightarrow V$ of singularities such that the induced rational map $\mu^{-1} \circ f \circ \mu: X \rightarrow V \rightarrow V \dashrightarrow X$ is a holomorphic étale endomorphism of X . This is known as the existence theorem of an equivariant resolution when f is an automorphism. However, the proof is also effective for non-isomorphic étale endomorphisms: A method of resolution of singularities is called to have a *functoriality* if, for any smooth morphism $X \rightarrow Y$, and for the resolutions of singularities $X' \rightarrow X$ and $Y' \rightarrow Y$ given by the method, X' is isomorphic to the fiber product $X \times_Y Y'$. The recent methods by Bierstone-Milman and by Villamayor using the notion of *invariant* have the functoriality (cf. [6], [14], [15], [55], [39]). Therefore, we call the resolution $X \rightarrow V$ above also an *equivariant resolution* even if f is a non-isomorphic étale endomorphism of V .

1.5. The meaning of our reduction. Let X be a nonsingular projective variety with an étale endomorphism f .

First, assume that X is uniruled. In view of Theorem C, f is considered to be built from a nearly étale endomorphism h of a non-uniruled normal variety Y up to isomorphism. Further, we can replace Y with a nonsingular variety and h with an étale endomorphism, provided that the minimal model conjecture is true for varieties birational to Y .

It is conjectured that a projective variety X is non-uniruled if and only if $\kappa(X) \geq 0$. This is regarded as a weak version of the abundance conjecture, and the three-dimensional case is proved by Miyaoka [43], which is a key to the proof of the three-dimensional abundance conjecture by [44] and [34]. The good minimal model conjecture is the combination of the minimal model conjecture and the abundance conjecture. Thus, under the assumption of good minimal model conjecture, the study of étale endomorphisms is reduced to that of étale endomorphisms of varieties of $\kappa \geq 0$.

Remark 1.1. A surjective endomorphism of a non-uniruled projective manifold X is étale. This follows from [30, Theorem 2] in the case of $\kappa(X) \geq 0$ (cf. [31, Theorem 11.7]). Fujimoto [20, Lemma 2.3] gives another proof in the same case. The proof also works in the case where $K_X H^{n-1} \geq 0$ for any ample divisor H of X ; this condition is satisfied if X is not uniruled by Miyaoka-Mori's criterion [45]. Thus, the étaleness for non-uniruled manifolds follows.

Second, assume that $\kappa(X) > 0$. Then we have the Iitaka fibration $\Phi: X \dashrightarrow Y$. By Theorem A, f induces a birational automorphism g of Y of finite order. Replacing X with a birational model, we may assume that Φ is holomorphic as remarked after Theorem A. Iterating f , we may assume f to be a morphism over Y , i.e., $\Phi \circ f = \Phi$. Then f induces an étale endomorphism of a general fiber F of Φ . This is a kind of reduction of f to an endomorphism of varieties of $\kappa = 0$. Theorem A reduces the dynamical study of holomorphic endomorphisms of compact Kähler manifolds of $\kappa > 0$ to the case of $\kappa = 0$, by the results in Appendix A. In fact, we show in Appendix A (cf. Theorem D) that $d_1(f) = d_1(f|_F)$ and $h_{\text{top}}(f) = h_{\text{top}}(f|_F)$ for the first dynamical degree d_1 and the topological entropy h_{top} .

However, even if we know the endomorphisms of fibers very well, it is rather difficult to determine the structure of f , as in the papers [20] and [21], which have determined the structure of endomorphisms of 3-dimensional projective manifolds of $\kappa \geq 0$.

Third, assume that $\kappa(X) = 0$. As in Section 1.2, an étale endomorphism f of X induces a nearly étale rational endomorphism of a weak Calabi-Yau variety F and an endomorphism of an abelian variety A , provided that the good minimal model conjecture is true. However, it is not clear that the nearly étale endomorphism induces an étale endomorphism of a certain resolution of singularities of F . Further, for the converse direction, it is not easy to recover the original endomorphism f from the two endomorphisms of F and A (cf. [20] for the 3-dimensional case).

Therefore, we can conclude, under the assumption of good minimal model conjecture, that the nearly étale endomorphisms of weak Calabi-Yau varieties and the endomorphisms

of abelian varieties are the building blocks of all the étale endomorphisms of complex projective manifolds.

For non-étale surjective endomorphisms (necessarily on uniruled manifolds), on the one hand, we know many examples of rationally connected varieties admitting non-isomorphic surjective endomorphisms, such as projective spaces, toric varieties, etc. (cf. [49]). On the other hand, at the moment, we do not have structure theorems for all the endomorphisms on them.

Organization of the paper. Section 2 is devoted to proving Theorem A and the application to the pluricanonical representation of the bimeromorphic automorphism group (cf. Corollary 2.4). In Section 3, we introduce a key notion of nearly étale map and study its elementary properties. Sections 4 and 5 are devoted to Theorem B and Theorem C, respectively. In Appendix A, we shall prove the equalities on the first dynamical degrees and the topological entropies mentioned above.

Notation and terminology. We refer to [35] for the definitions of minimal models and of singularities including terminal, canonical, log-terminal, and rational singularities. Also, we refer to [38], [41], [46] for additional information on the birational geometry and the minimal model theory. Here we add an explanation of the notion of *fibration* (or *fiber space*): Let $\pi: X \rightarrow Y$ be a proper morphism of normal algebraic (resp. complex analytic) varieties. If a general fiber of π is connected, then every fiber is connected and $\mathcal{O}_Y \simeq \pi_* \mathcal{O}_X$. In this case, π is called a fibration or a fiber space. A proper rational map (cf. Definition 3.1 below) (resp. a proper meromorphic map) between normal varieties is called a fibration or a fiber space if it is the composite of a proper birational (resp. bimeromorphic) map and a holomorphic fibration.

Acknowledgement. The first named author expresses his gratitude to the Department of Mathematics of the National University of Singapore for the hospitality during his stay in October 2006. The work of this paper is based on the discussion with the second named author during the stay. The authors are grateful to Dr. Hiraku Kawanoue for answering questions on equivariant resolutions, and to Professor Takeshi Abe for pointing out a gap in the proof of Proposition 2.3 in an earlier version. The authors would like to thank Professor Yoshio Fujimoto for his encouragement and the referee for the suggestions.

2. THE CASE OF POSITIVE KODAIRA DIMENSION

2.1. Iitaka fibration. In the situation of Theorem A, we may assume that X is a compact Kähler manifold, without loss of generality. We have a natural isomorphism $f^*: H^0(X, mK_X) \xrightarrow{\sim} H^0(X, mK_X)$. In fact, there is a bimeromorphic morphism $\mu: Z \rightarrow$

X from another compact Kähler manifold Z such that $\varphi := f \circ \mu: Z \rightarrow X$ is holomorphic; then f^* is expressed as the composite

$$H^0(X, mK_X) \xrightarrow{\varphi^*} H^0(Z, mK_Z) \xrightarrow[\simeq]{(\mu^*)^{-1}} H^0(X, mK_X),$$

which does not depend on the choice of $\mu: Z \rightarrow X$. The pullback f^* induces an automorphism g of $|mK_X|^\vee$ preserving W_m . The problem is to show the finiteness of the order of $g \in \text{Aut}(W_m)$. We begin with the following simple result.

Lemma 2.1. *If Theorem A does not hold, then there is a positive-dimensional connected commutative linear algebraic subgroup $G \subset \text{Aut}(W_m)$ such that $g^k \in G$ for some $k > 0$. In particular, W_m is ruled in this case.*

Proof. Let $\tilde{G} \subset \text{PGL} = \text{Aut}(|mK_X|^\vee)$ be the Zariski closure of the cyclic group $I = \{g^j \mid j \in \mathbb{Z}\}$. Then $\tilde{G}$ is abelian, since it is contained in the abelian algebraic subgroup $Z(I) \cap Z(Z(I))$, where $Z(S)$ denotes the algebraic subgroup $\{\gamma \in \text{PGL} \mid \gamma s = s\gamma \text{ for any } s \in S\}$ for a subset $S \subset \text{PGL}$. Let G be the identity connected component of $\tilde{G}$. Then $\dim G > 0$ and $g^k \in G$ for some $k > 0$. Since the action of G preserves W_m , G acts non-trivially on W_m . Thus W_m is ruled by a result of Matsumura (cf. [52, Theorem 14.1]). $\square$

Remark 2.2. There is another proof of Lemma 2.1 by an argument similar to [52, Proposition 14.7]: In fact, we can show that f^* is expressed as a diagonal matrix. Thus, G is contained in an algebraic torus. We can prove more on f^* by the argument of [52, Proposition 14.4]: If λ is an eigenvalue of $f^*: H^0(X, mK_X) \rightarrow H^0(X, mK_X)$, then λ is an algebraic integer with $|\lambda|^{2/m} = \deg f$.

The following is a key to the proof of Theorem A.

Proposition 2.3. *Let $\pi: X \rightarrow Y$ be a fiber space of a compact Kähler manifold X over a nonsingular rational curve $Y \simeq \mathbb{P}^1$. Let $f: X \dashrightarrow X$ be a dominant meromorphic map and $g: Y \xrightarrow{\simeq} Y$ an automorphism such that $\pi \circ f = g \circ \pi$. If $\kappa(X) \geq 0$, then g is of finite order.*

Proof. Step 1. We first prove in the case where $p_g(X) = \dim H^0(X, K_X) > 0$. Since π is smooth over a dense open subset U of Y , we have a variation of Hodge structure $H_U = R^d \pi_* \mathbb{Z}_X|_U$ for $d = \dim X/Y = \dim X - 1$.

Let $\mu: Z \rightarrow X$ be a bimeromorphic morphism from another compact Kähler manifold Z such that $\varphi := f \circ \mu: Z \rightarrow X$ is holomorphic. Then

$$f_\omega^*: g^*(\pi_* \omega_{X/Y}) \xrightarrow{\varphi^*} g^*(\pi_* \varphi_* \omega_{Z/Y}) \xrightarrow[\simeq]{(\mu^*)^{-1}} \pi_* \omega_{X/Y}$$

is injective. Note that $\pi_* \omega_{X/Y}$ is just the d -th filter $\mathcal{F}^d({}^u\mathcal{H})$ of the upper canonical extension ${}^u\mathcal{H}$ of $H_U \otimes \mathcal{O}_U$ in the sense of Kollár [36].

We have also the pullback homomorphism

$$f_{\text{coh}}^*: g^{-1}(\mathbf{R}^d \pi_* \mathbb{Z}_X) \xrightarrow{\varphi^*} g^{-1} \mathbf{R}^d (\pi \circ \varphi)_* \mathbb{Z}_Z \xrightarrow{\mu_*} \mathbf{R}^d \pi_* \mathbb{Z}_X,$$

where μ_* is induced from the trace map $\mathbf{R} \mu_* \mathbb{Z}_Z[2n] \rightarrow \mathbb{Z}_X[2n]$ for $n = \dim X$. Note that f_{coh}^* is compatible with f_ω^* , i.e., the diagram

$$\begin{array}{ccc} (g^{-1} \mathbf{R}^d \pi_* \mathbb{Z}_X)|_{U'} \otimes \mathcal{O}_{U'} & \xrightarrow{f_{\text{coh}}^*} & (\mathbf{R}^d \pi_* \mathbb{Z}_X)|_{U'} \otimes \mathcal{O}_{U'} \\ \uparrow & & \uparrow \\ (g^* \pi_* \omega_{X/Y})|_{U'} & \xrightarrow{f_\omega^*} & \pi_* \omega_{X/Y}|_{U'} \end{array}$$

is commutative over the open subset $U' = U \cap g^{-1}U$. Let $J_k \subset \mathbf{R}^d \pi_* \mathbb{Z}_X$ be the image of

$$\begin{aligned} (f_{\text{coh}}^*)^k: (g^k)^{-1}(\mathbf{R}^d \pi_* \mathbb{Z}_X) &\xrightarrow{(g^{k-1})^{-1} f_{\text{coh}}^*} (g^{k-1})^{-1}(\mathbf{R}^d \pi_* \mathbb{Z}_X) \rightarrow \cdots \\ &\cdots \rightarrow g^{-1}(\mathbf{R}^d \pi_* \mathbb{Z}_X) \xrightarrow{f_{\text{coh}}^*} \mathbf{R}^d \pi_* \mathbb{Z}_X \end{aligned}$$

for $k > 0$. Then $J_k \supset J_{k+1}$ and $f_{\text{coh}}^*(g^{-1}J_k) = J_{k+1}$. Moreover, for $k \gg 0$, $f_{\text{coh}}^*: g^{-1}J_k \rightarrow J_{k+1}$ is isomorphic and $J_k \otimes \mathbb{Q} = J_{k+1} \otimes \mathbb{Q}$, since any stalk of $\mathbf{R}^d \pi_* \mathbb{Z}_X$ is a finitely generated abelian group. We set $J := J_k$ for certain $k \gg 0$. Then, for a non-empty Zariski open subset $U'' \subset U$, $J|_{U''}$ defines a variation of Hodge substructure of $\mathbf{R}^d \pi_* \mathbb{Z}_X|_{U''}$, and $J|_{U''} \otimes \mathcal{O}_{U''}$ contains $\pi_* \omega_{X/Y}|_{U''}$ as the d -th Hodge filter. Furthermore, the injection $f_{\text{coh}}^*: g^{-1}J \hookrightarrow J$ is compatible with f_ω^* . Let U_{max} be the maximal open subset of Y such that there is a variation of Hodge structure J_{max} on U_{max} with $J_{\text{max}}|_{U''} \simeq J|_{U''}$. Then $g^{-1}U_{\text{max}} = U_{\text{max}}$. Thus, we may assume that $Y \setminus U_{\text{max}}$ consists of at most two points; otherwise, g is of finite order. Note that a Kähler form of X defines a real polarization of the variation of Hodge structure J_{max} .

Suppose that $U_{\text{max}} = Y \simeq \mathbb{P}^1$ or $U_{\text{max}} \simeq \mathbb{C}$. Then U_{max} is simply connected and we have a period map from U_{max} to the period domain associated with the polarized variation of Hodge structure J_{max} (we need the polarization here). Griffiths proves that this map is “horizontal” (cf. [23, Sections 8 and 9]) and satisfies a kind of Schwarz inequality ([23, Theorem 10.1]). Thus, this is a constant map by a reason similar to the fact that the map from $\mathbb{C}$ to a Kobayashi hyperbolic manifold is constant; in this case, the monodromy representation is also trivial, so the variation of Hodge structure J_{max} is constant, i.e., the local system is trivial and every Hodge filtration $F^p(J_{\text{max}})$ is a trivial locally free sheaf. Hence, the sheaf $\pi_* \omega_{X/Y} \simeq F^d(J_{\text{max}})$ is trivial. This leads to a contradiction: $H^0(X, K_X) \simeq H^0(Y, \pi_*(\omega_X)) = H^0(Y, \pi_*(\omega_{X/Y}) \otimes \omega_Y) = H^0(\mathbb{P}^1, \mathcal{O}(-2)) = 0$.

Hence, we may assume $U_{\text{max}} = \mathbb{C}^*$. Then, the period map associated with J_{max} is constant, since the universal covering space of U_{max} is $\mathbb{C}$. In particular, the image of the monodromy representation $\pi_1(U_{\text{max}}, y) \rightarrow \text{Aut}(J_{\text{max}, y})$ is finite for a reference point $y \in Y$.

Let $\tau: Y' \simeq \mathbb{P}^1 \rightarrow Y$ be a cyclic covering étale over $U_{\max}$ such that $\tau^{-1}J_{\max}$ extends to a trivial constant sheaf of Y' . We may assume that g lifts to an automorphism g' of Y' . Let $X' \rightarrow X \times_Y Y'$ be a resolution of singularities and let $\pi': X' \rightarrow Y'$ be the induced morphism. Then $f \times g'$ induces a meromorphic endomorphism of X' and $p_g(X') > 0$. Since $\tau^{-1}J_{\max}$ has trivial monodromy, a similar variation of Hodge structure $J'_{\max}$ is defined on Y' . Thus $\pi'_*\omega_{X'/Y'}$ is a free $\mathcal{O}_{Y'}$ -module, and we have the same contradiction as above.

Step 2. General case: Let $s \in H^0(X, mK_X)$ be an eigenvector of f^* . We shall consider a cyclic covering corresponding to taking the m -th root of s : Let $\mathcal{A} = \bigoplus_{i=0}^{m-1} \mathcal{O}_X(-iK_X)$ be the $\mathcal{O}_X$ -algebra determined by $s: \mathcal{O}_X(-mK_X) \rightarrow \mathcal{O}_X$. Taking a resolution $\widehat{X}$ of singularities of $\text{Spec}_X \mathcal{A}$, let $\tau: \widehat{X} \rightarrow X$ be the composite $\widehat{X} \rightarrow \text{Spec}_X \mathcal{A} \rightarrow X$. Then $\tau^*s \in H^0(\widehat{X}, mK_{\widehat{X}})$ is expressed as σ^m for a section $\sigma \in H^0(\widehat{X}, K_{\widehat{X}})$. Let X' be a connected component of $\widehat{X}$. Then $\kappa(X') = \kappa(X)$ and $p_g(X') > 0$. Let $\pi': X' \rightarrow Y'$ be the fiber space and let $\lambda: Y' \rightarrow Y$ be the finite morphism obtained as the Stein factorization of $X' \rightarrow X \rightarrow Y$.

Since s is an eigenvector of f^* , we have a meromorphic map $\hat{f}: \widehat{X} \cdots \rightarrow \widehat{X}$ with $\tau \circ \hat{f} = f \circ \tau$. Replacing f with a suitable power f^k , we may assume that $\hat{f}$ preserves X' . Let $f': X' \cdots \rightarrow X'$ be the induced rational map. Then there is an automorphism g' of Y' such that $\pi' \circ f' = g' \circ \pi'$ and $\lambda \circ g' = g \circ \lambda$. If g' is of finite order, then so is g . If the genus of Y' is greater than one, then g' is of finite order. Even if the genus of Y' is one, g' is of finite order since g' preserves the ample invertible sheaf $\lambda^*\mathcal{O}(1)$. If the genus of Y' is zero, then g' is of finite order by *Step 1*. Thus, we are done. $\square$

Remark. If f is a holomorphic endomorphism of X and if X is projective, then Proposition 2.3 follows from [54], since π has at least three singular fibers preserved by g .

2.2. Proof of Theorem A. We first compare W_m and W_{ml} for positive integers m and l with $|mK_X| \neq \emptyset$. There is a natural rational map $\Psi_{m,ml}: W_{ml} \cdots \rightarrow W_m$ such that $\Phi_m = \Psi_{m,ml} \circ \Phi_{ml}$. In fact, $\Psi_{m,ml}$ is defined by the commutative diagram

$$\begin{array}{ccc} W_{ml} & \hookrightarrow & |mlK_X|^\vee = \mathbb{P}(H^0(X, mlK_X)) \\ \Psi_{m,ml} \downarrow & & \downarrow \mu \\ W_m & \hookrightarrow & |mK_X|^\vee \xhookrightarrow{\iota} \mathbb{P}(\text{Sym}^l H^0(X, mK_X)), \end{array}$$

where ι is the Veronese embedding and μ is induced from the natural homomorphism $\text{Sym}^l H^0(X, mK_X) \rightarrow H^0(X, mlK_X)$. Let g_m and g_{ml} , respectively, be the automorphisms of W_m and W_{ml} induced from f . Then $g_m \circ \Psi_{m,ml} = \Psi_{m,ml} \circ g_{ml}$ by construction. Thus, we may replace m with any multiple ml in order to prove Theorem A. Therefore, we may assume from the beginning that $\Phi_m: X \cdots \rightarrow W_m$ gives rise to the Iitaka fibration of X .

We shall derive a contradiction by assuming that $g := g_m$ is of infinite order. By Lemma 2.1, we may assume that g is contained in a positive-dimensional connected commutative linear algebraic group $G \subset \text{Aut}(W_m)$. Let $Y \rightarrow W_m$ be an equivariant resolution of singularities so that G acts on Y holomorphically. There is a sequence

$$\{\text{id}\} = G_0 \subset G_1 \subset G_2 \subset \cdots \subset G_l = G$$

of algebraic subgroups such that $\dim G_i/G_{i-1} = 1$ for $1 \leq i \leq l$. In particular, $G_i/G_{i-1} \simeq \mathbb{G}_m$ or $\mathbb{G}_a$. Let $Y \cdots \rightarrow Y_1 \subset \text{Hilb}(Y)$ be the meromorphic quotient of Y by G_1 (cf. [19, Theorem 4.1]). Then G/G_1 acts on Y_1 . Replacing Y and Y_1 by their nonsingular models, respectively, we may assume that $Y \rightarrow Y_1$ is holomorphic and G -equivariant. Let $Y_1 \cdots \rightarrow Y_2 \subset \text{Hilb}(Y_1)$ be the meromorphic quotient of Y_1 by G_2/G_1 , and replace Y , Y_1 , and Y_2 by their nonsingular models, respectively, so that $Y \rightarrow Y_1$ and $Y_1 \rightarrow Y_2$ are G -equivariant morphisms. Continuing similar constructions, we have a sequence of G -equivariant morphisms

$$Y = Y_0 \rightarrow Y_1 \rightarrow \cdots \rightarrow Y_l$$

such that $Y \rightarrow Y_i$ is birational to the meromorphic quotient by G_i . Then, for $0 \leq i < l$, a general fiber of $Y_i \rightarrow Y_{i+1}$ is a smooth rational curve which is the closure of an orbit of G_{i+1}/G_i . There is a similar and stronger assertion in [42, Theorem 4.6].

Replacing X by its blowup, we may assume that the composite $X \cdots \rightarrow W_m \cdots \rightarrow Y$ is holomorphic. Let us consider the composition $\phi_i: X \rightarrow Y \rightarrow Y_i$ for $1 \leq i \leq l$. For a very general point $y_i \in Y_i$, let $F_i \subset X$ be the fiber $\phi_i^{-1}(y_i)$ and let $C_{i-1} \subset Y_{i-1}$ be the fiber of $Y_{i-1} \rightarrow Y_i$ over y_i . Then $\kappa(F_i) > 0$ for $i > 0$ by the easy addition formula:

$$\kappa(X) = \dim Y \leq \kappa(F_i) + \dim Y_i.$$

Now g acts on Y_l trivially. Thus $f: X \cdots \rightarrow X$ is a meromorphic map over Y_l , and a dominant meromorphic map $F_l \cdots \rightarrow F_l$ is induced. By Proposition 2.3, the action of g on C_{l-1} is of finite order. Thus g^k acts on Y_{l-1} trivially for some $k > 0$. Hence, $\{g^{kj} \mid j \in \mathbb{Z}\} \subset G_{l-1}$. This contradicts that G is the identity component of the Zariski closure of $\{g^j \mid j \in \mathbb{Z}\}$. This completes the proof of Theorem A. $\square$

Theorem A has an application to the pluricanonical representations of the bimeromorphic automorphism group $\text{Bim}(X)$ of compact complex manifolds X in the class $\mathcal{C}$. The following result is proved in [52, §14] (cf. [47]) when X is a Moishezon manifold.

Corollary 2.4. *Let X be a compact complex manifold in the class $\mathcal{C}$ and let*

$$\rho_m: \text{Bim}(X) \rightarrow \text{Aut}(H^0(X, mK_X))$$

be the m -th pluricanonical representation (cf. [52, §14]) for a positive integer m with $H^0(X, mK_X) \neq 0$. Then the image of the induced group homomorphism below is finite:

$$\rho'_m: \text{Bim}(X) \rightarrow \text{PGL}(H^0(X, mK_X)) = \text{Aut}(H^0(X, mK_X))/\mathbb{C}^\star.$$

Proof. Let W_m be the image of the m -th pluricanonical map $\Phi_m: X \dashrightarrow |mK_X|^\vee$. Then W_m is not contained in any hyperplane of $|mK_X|^\vee$. Thus, for $\gamma \in \text{Bim}(X)$, the order of $\rho'_m(\gamma)$ equals the order of the action of γ on W_m . By Theorem A, the order of $\rho'_m(\gamma)$ is finite for any $\gamma \in \text{Bim}(X)$. So $\rho_m(\gamma)$ is expressed as a diagonal matrix and its eigenvalues are $(\alpha\theta_1, \alpha\theta_2, \dots, \alpha\theta_k)$ for $k = \dim H^0(X, mK_X)$, for a constant $\alpha \in \mathbb{C}^\star$, and for roots θ_i of unity, where $\theta_1 = 1$. By [52, Proposition 14.4] and by an argument in [52, Theorem 14.10], $\alpha\theta_i$ are algebraic integers, $|\alpha| = 1$, and the degree $[\mathbb{Q}(\alpha\theta_i) : \mathbb{Q}]$ of field extension $\mathbb{Q}(\alpha\theta_i)/\mathbb{Q}$ is bounded above by a suitable constant N which depends neither on i nor on γ . Thus the degree $[\mathbb{Q}(\theta_i) : \mathbb{Q}]$ is bounded above by N^2 for any i . Hence, the order of $\rho'_m(\gamma)$ is uniformly bounded. Therefore, the image of ρ'_m is a finite group by a theorem of Burnside (cf. [52, Theorem 14.9]). $\square$

3. NEARLY ÉTALE MAPS

We introduce the notion of nearly étale map and study its basic properties.

Definition 3.1 (cf. [31]). Let $h: V \dashrightarrow W$ be a rational (resp. meromorphic) map between algebraic (resp. complex analytic) varieties. The map h is called *proper* if the projections $p_1: \Gamma_h \rightarrow V$ and $p_2: \Gamma_h \rightarrow W$ are both proper for the graph $\Gamma_h \subset V \times W$. For algebraic varieties V and W , V is called *proper birational* to W if there exists a proper birational map $V \dashrightarrow W$.

Remark. The first projection $p_1: \Gamma_h \rightarrow V$ is proper for any meromorphic map h . In particular, a bimeromorphic map is always proper.

We extend the notion of Stein factorization to the case of proper rational (resp. proper meromorphic) maps.

Definition 3.2. Let $h: V \dashrightarrow W$ be a proper rational (resp. proper meromorphic) map from a normal variety V to an algebraic (resp. complex analytic) variety W . Then there exists a finite morphism $\tau: W^\sharp \rightarrow W$ uniquely up to isomorphism over W such that $W^\sharp$ is normal and $h = \tau \circ \varphi$ for a rational (resp. meromorphic) fiber space $\varphi: V \dashrightarrow W^\sharp$. The factorization $V \dashrightarrow W^\sharp \rightarrow W$ of h is called the *Stein factorization*.

Remark. If h is holomorphic, then this is the usual Stein factorization, where τ is determined by the isomorphism $h_*\mathcal{O}_V \simeq \tau_*\mathcal{O}_{W^\sharp}$. In case h is not holomorphic, let $\mu: V' \rightarrow V$ be a proper birational (resp. bimeromorphic) morphism such that V' is normal and

$h' = h \circ \mu: V' \rightarrow W$ is holomorphic; then τ is just given by the Stein factorization of h' . The uniqueness of τ in this case follows from Zariski's main theorem. If V and W are algebraic, then $W^\sharp$ is just the normalization of W in the function field $\mathbb{C}(V)$.

Definition 3.3. Let $h: V \dashrightarrow W$ be a proper rational (resp. meromorphic) map between algebraic (resp. complex analytic) varieties. The map h is called *nearly étale* if there exist proper birational (resp. bimeromorphic) maps $\mu: Y \dashrightarrow W$, $\nu: X \dashrightarrow V$ and a morphism $f: X \rightarrow Y$ such that

- (1) X and Y are algebraic (resp. complex analytic) varieties,
- (2) f is a finite étale morphism, and
- (3) $\mu \circ f = h \circ \nu$, i.e., the diagram below is commutative:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \nu \downarrow & & \downarrow \mu \\ V & \xrightarrow{h} & W. \end{array}$$

Remark.

- A nearly étale map is dominant and generically finite.
- A proper birational (resp. bimeromorphic) map is nearly étale.
- A finite étale morphism is nearly étale.
- Any nearly étale rational (resp. meromorphic) map $V \dashrightarrow W$ is the composition of a proper birational (resp. bimeromorphic) map $V \dashrightarrow W^\sharp$ and a nearly étale finite morphism $W^\sharp \rightarrow W$. In fact, it is enough to take the Stein factorization of the rational (resp. meromorphic) map from the normalization of V to W .

We now study basic properties of nearly étale maps.

Lemma 3.4. *Let $h: V \dashrightarrow W$ be a nearly étale rational (resp. meromorphic) map between normal varieties. Suppose that $\pi_1^{\text{alg}}(W) \simeq \pi_1^{\text{alg}}(Y)$ for a resolution $Y \rightarrow W$ of singularities of W , where π_1^{alg} stands for the algebraic fundamental group. Then the finite morphism $W^\sharp \rightarrow W$ is étale for the Stein factorization $V \dashrightarrow W^\sharp \rightarrow W$ of h .*

Proof. The fundamental group π_1 is a proper birational (resp. bimeromorphic) invariant for nonsingular varieties. Therefore, there is a finite étale covering $W' \rightarrow W$ corresponding to the finite-index subgroup $\pi_1(X) \subset \pi_1(Y)$ via the isomorphism $\pi_1(Y) \simeq \pi_1(W)$ for the étale covering $X \rightarrow Y$ in Definition 3.3. Since X is proper birational (resp. bimeromorphic) to W' , we have $W' \simeq W^\sharp$ over W . $\square$

As a consequence of Lemma 3.4, we can show that the composition of two nearly étale maps are also nearly étale. Our next result shows that a nearly étale finite morphism is turned to be an étale morphism by a suitable base change.

Corollary 3.5. *Let $V \rightarrow W$ be a nearly étale finite morphism between normal varieties. Let $Z \rightarrow W$ be a morphism from a normal variety Z such that the image contains a non-empty open subset of W and $\pi_1^{\text{alg}}(Z) \simeq \pi_1^{\text{alg}}(M)$ for a resolution $M \rightarrow Z$ of singularities of Z . Let $\tilde{V}_Z$ be the normalization of $V \times_W Z$ and $\tilde{V}_Z \rightarrow Z^\sharp \rightarrow Z$ the Stein factorization of the morphism induced from the second projection $V \times_W Z \rightarrow Z$. Then $Z^\sharp \rightarrow Z$ is étale.*

Proof. Let $\mu: Y \rightarrow W$ be a proper birational (resp. bimeromorphic) morphism from a nonsingular variety Y . By Lemma 3.4, we have a finite étale covering $X \rightarrow Y$ such that $V \rightarrow W$ is obtained as the Stein factorization of $X \rightarrow Y \rightarrow W$. There is a morphism $Y' \rightarrow Y \times_W Z$ from a nonsingular variety Y' such that the composite $Y' \rightarrow Y \times_W Z \rightarrow Z$ is proper birational (resp. bimeromorphic), since the image of $Z \rightarrow W$ contains a non-empty open subset. Let $X' \rightarrow Y'$ be the finite étale morphism obtained as the pullback of $X \rightarrow Y$ by $Y' \rightarrow Y \times_W Z \rightarrow Y$. We infer that any connected component of $Z^\sharp$ is obtained as the Stein factorization of the morphism $X' \rightarrow Y' \rightarrow Z$ restricted to a connected component of X' . Therefore, $Z^\sharp \rightarrow Z$ is étale by Lemma 3.4. $\square$

Definition 3.6 (cf. [37, (7.1.2)], [48, Section 2]). Let (V, P) be a germ of normal complex analytic singularity and $\mu: Z \rightarrow V$ a resolution of singularity.

- (V, P) is called an *algebraically π_1 -rational singularity* if the algebraic fundamental group of $\mu^{-1}(P)$ is trivial.
- (V, P) is a *π_1 -rational singularity* if the fundamental group of $\mu^{-1}(P)$ is trivial.

Let W be a normal variety.

- W_{reg} denotes the nonsingular locus of W .
- W_{rat} denotes the set of points $P \in W$ such that (W, P) is nonsingular or is a rational singularity.
- W_{apr} denotes the set of points $P \in W$ such that (W, P) is an algebraically π_1 -rational singularity.

Remark 3.7.

- (1) If (V, Δ) is log-terminal (klt) at the point P for a $\mathbb{Q}$ -divisor Δ with $\lfloor \Delta \rfloor = 0$, then (V, P) is an algebraically π_1 -rational singularity by [37, Theorem 7.5], and is in fact a π_1 -rational singularity by [51].
- (2) W_{reg} and W_{rat} are Zariski open dense subset of a normal variety W , but W_{apr} is not necessarily open.

- (3) If $W = W_{\text{apr}}$, then $\pi_1^{\text{alg}}(W) \simeq \pi_1^{\text{alg}}(W)$ for any resolution $Y \rightarrow V$ of singularities of W . In particular, this holds if (W, Δ) is log-terminal for a $\mathbb{Q}$ -divisor Δ with $\lfloor \Delta \rfloor = 0$ (cf. [37, Theorem 7.8]).

The result below says that a nearly étale endomorphism induces an étale endomorphism of a certain nonsingular model, provided that the minimal model conjecture is true.

Lemma 3.8. *Let $h: V \rightarrow V$ be a nearly étale finite morphism for a normal variety V . Assume the existence of the relative canonical model V_{can} for the resolutions of singularities of V ; equivalently, assume that the sheaf*

$$\mathcal{R}_V := \bigoplus_{m \geq 0} \mu_* \mathcal{O}_M(mK_M)$$

of relative canonical ring is a finitely generated $\mathcal{O}_V$ -algebra for a resolution $\mu: M \rightarrow V$ of singularities of V , where $V_{\text{can}} = \text{Proj}_V \mathcal{R}_V$. Then h lifts to an étale endomorphism of V_{can} and to an étale endomorphism of a certain resolution of singularities of V .

Proof. Since V_{can} has only π_1 -rational singularities, applying Corollary 3.5 to $V_{\text{can}} \rightarrow V$, we have a finite étale morphism $h_{\text{can}}: V_{\text{can}}^\# \rightarrow V_{\text{can}}$ and a proper birational (resp. bimeromorphic) morphism $V_{\text{can}}^\# \rightarrow V$ such that

$$\begin{array}{ccc} V_{\text{can}}^\# & \longrightarrow & V \\ h_{\text{can}} \downarrow & & \downarrow h \\ V_{\text{can}} & \longrightarrow & V \end{array}$$

is commutative. Then $V_{\text{can}}^\#$ is also the relative canonical model for the resolutions of singularities of V , since it has only canonical singularities and its canonical divisor is also relatively ample over V . So $V_{\text{can}}^\# \simeq V_{\text{can}}$ and h_{can} is regarded as an étale endomorphism of V_{can} . Applying the equivariant resolution of singularities of V_{can} with respect to the étale endomorphism h_{can} in the sense of Section 1.4, we have a lift to a certain resolution of singularities of V . $\square$

Lemma 3.9 (cf. [37, Theorem 5.2]). *For a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \psi \downarrow & & \downarrow \pi \\ V & \xrightarrow{h} & W \end{array}$$

of proper surjective morphisms of normal complex analytic varieties, h is étale provided that the following conditions are satisfied:

- (1) X and Y are nonsingular.
- (2) ψ and π have connected fibers.

- (3) *The induced morphism $X \rightarrow V \times_W Y$ is an isomorphism over a dense open subset of V .*
- (4) *f is a finite étale morphism, and h is a finite morphism.*
- (5) *$R^i \pi_* \mathcal{O}_Y = 0$ and $R^i \psi_* \mathcal{O}_X = 0$ for any $i > 0$.*

Proof. We may assume that $\deg f = \deg h > 1$. Let $P \in W$ be an arbitrary point. It is enough to show that the cardinality $\sharp h^{-1}(P)$ of $h^{-1}(P)$ equals $\deg h$. For the proof, we may replace Y with another nonsingular variety Y' by taking a bimeromorphic morphism $Y' \rightarrow Y$. In fact, the pullback $f': X' \rightarrow Y'$ of $f: X \rightarrow Y$ by $Y' \rightarrow Y$ induces a similar commutative diagram satisfying the same conditions. Thus, we may assume that the reduced structure $E = \pi^{-1}(P)_{\text{red}}$ of the fiber $\pi^{-1}(P) = Y \times_W \{P\}$ is a normal crossing divisor, after blowing up $\pi^{-1}(P)$. Then $f^{-1}E$ is also a reduced normal crossing divisor of X . Here, $f^{-1}E$ is the disjoint union of the reduced structures $\tilde{E}_Q$ of the fibers $\psi^{-1}(Q)$ for points $Q \in h^{-1}(P)$.

We shall show $\chi(E, \mathcal{O}_E) = 1$. In the natural commutative diagram

$$\begin{array}{ccc} (R^i \pi_* \mathbb{C}_Y)_P & \xrightarrow{\cong} & H^i(E, \mathbb{C}) \\ \downarrow & & \downarrow \\ (R^i \pi_* \mathcal{O}_Y)_P & \longrightarrow & H^i(E, \mathcal{O}_E), \end{array}$$

the right vertical arrow is surjective by the theory of mixed Hodge structures on normal crossing varieties. Therefore, $H^i(E, \mathcal{O}_E) = 0$ for $i > 0$ by the condition (5). In particular, $\chi(E, \mathcal{O}_E) = 1$. By the same argument, we also have $\chi(\tilde{E}_Q, \mathcal{O}) = 1$. On the other hand, $\chi(f^{-1}E, \mathcal{O}) = (\deg f)\chi(E, \mathcal{O}_E) = \deg f$, since f is étale. Therefore,

$$\sharp h^{-1}(P) = \sum \chi(\tilde{E}_Q, \mathcal{O}) = \chi(f^{-1}E, \mathcal{O}) = \deg f = \deg h. \quad \square$$

Proposition 3.10. *Let $h: V \rightarrow W$ be a nearly étale finite morphism between normal varieties. Then:*

- (1) *$h^{-1}(W_{\text{apr}}) \subset V_{\text{apr}}$ and h is étale along $h^{-1}(W_{\text{apr}})$.*
- (2) *h is étale along V_{rat} .*
- (3) *$h^{-1}(W_{\text{apr}} \cap W_{\text{rat}}) = V_{\text{apr}} \cap V_{\text{rat}}$.*
- (4) *$h^{-1}(W_{\text{reg}}) = V_{\text{reg}}$.*

Proof. (1) follows from Lemma 3.4 applied to an open neighborhood of a point in W_{apr} . Moreover, (3) and (4) are derived from (1) and (2), since $h(V_{\text{rat}}) \subset W_{\text{rat}}$. Therefore, it suffices to show (2).

Let P be a point of V_{rat} . In order to prove the étaleness of h at P , we may replace W with an open neighborhood $\mathcal{W}$ of $h(P)$ and V with a connected component of $h^{-1}(\mathcal{W})$ containing P . Thus, we may assume that W is simply connected and $h^{-1}(h(P)) = \{P\}$,

set-theoretically. Let $\mu: Y \rightarrow W$ be a resolution of singularities. Then, by Lemma 3.4, we have a finite étale covering $f: X \rightarrow Y$ and a bimeromorphic morphism $\nu: X \rightarrow V$ such that $\mu \circ f = h \circ \nu$. Applying Lemma 3.9 to the commutative diagram expressing $\mu \circ f = h \circ \nu$, we infer that h is an isomorphism. Thus, we are done. $\square$

For a nearly étale endomorphism of a normal algebraic variety, we have the following additional property:

Proposition 3.11. *Let $h: V \rightarrow V$ be a nearly étale finite surjective endomorphism of a normal algebraic variety V . Then $h^{-1}(V_{\text{rat}}) = V_{\text{rat}}$.*

Proof. Let B be the complement of V_{rat} in V . Then we have the inclusion $h^{-1}(B) \subset B$ by Proposition 3.10. The other inclusion follows from Lemma 3.12 below. $\square$

Lemma 3.12. *Let $h: V \rightarrow V$ be a finite surjective endomorphism of an algebraic variety V . Suppose $h^{-1}(B) \subset B$ for a closed subset $B \subset V$. Then $h^{-1}(B) = B$. Further, $\Gamma \mapsto h(\Gamma)$ induces an automorphism of the set I_B of irreducible components Γ of B . The inverse map is given by $\Gamma \mapsto h^{-1}(\Gamma)$.*

Proof. Let $I_B^{(k)}$ be the set of irreducible components of B of dimension k . Then $I_B = \bigcup I_B^{(k)}$. Let $J_B^{(k)}$ be the subset of $I_B^{(k)}$ consisting of $\Gamma \in I_B$ with $h(\Gamma) \subset B$. It is enough to show the following two properties for any k :

- (1)_k: $I_B^{(k)} = J_B^{(k)}$.
- (2)_k: $h_*: J_B^{(k)} \rightarrow I_B^{(k)}$ given by $\Gamma \mapsto h(\Gamma)$ is bijective.

We shall prove by descending induction on k . We set $d = \dim B$. For $\Gamma \in I_B^{(d)}$, some $\Gamma' \in I_B^{(d)}$ is contained in $h^{-1}(\Gamma)$. Here, $\Gamma' \in J_B^{(d)}$, since $h(\Gamma') = \Gamma$. Hence $h_*: J_B^{(d)} \rightarrow I_B^{(d)}$ is surjective. Since $I_B^{(d)}$ is a finite set, $J_B^{(d)} = I_B^{(d)}$ and $h_*: J_B^{(d)} \rightarrow I_B^{(d)}$ is bijective.

Next, assume that (1)_l and (2)_l hold for any integer $l > k$. If $\Gamma \in I_B^{(k)}$, then an irreducible component Δ of $h^{-1}(\Gamma)$ dominates Γ , and $\Delta \subset \Gamma'$ for some $\Gamma' \in I_B$. If $\dim \Gamma' > k$, then $h(\Gamma') \subset B$ by induction, and hence $\Gamma \subset h(\Gamma')$; this is a contradiction. Thus $\dim \Gamma' = k$, $\Gamma' = \Delta$, and $\Gamma' \in J_B^{(k)}$. Hence, $h_*: J_B^{(k)} \rightarrow I_B^{(k)}$ is surjective. Therefore, $I_B^{(k)} = J_B^{(k)}$ and $h_*: J_B^{(k)} \rightarrow I_B^{(k)}$ is bijective. Thus, we are done. $\square$

If $f: X \rightarrow X$ is a surjective endomorphism of a nonsingular projective variety X with $\kappa(X) \geq 0$, then f is étale. But if we drop the condition of nonsingularity, then we can expect neither the étaleness nor even the nearly étaleness. Indeed, we have:

Example 3.13. Let A be an abelian surface and V the quotient space of A by the involution $\iota: A \ni a \mapsto -a \in A$ with respect to a group structure of A . The minimal resolution of singularities of V is a K3 surface called the Kummer surface associated with A and is

denoted by $\mathrm{Km}(A)$. The variety V has only canonical singularities and $K_V \sim 0$. Let $\mu = \mu_m: A \rightarrow A$ be the multiplication map $a \mapsto ma$ by an integer m with $|m| > 1$. Then it descends to a non-isomorphic surjective endomorphism $\mu_V: V \rightarrow V$. Here, μ_V is not nearly étale since $\mathrm{Km}(A)$ is simply connected.

4. THE CASE OF ZERO KODAIRA DIMENSION

4.1. Albanese closure.

Definition 4.1. For a normal projective variety V , we define $q^{\max}(V)$ to be the supremum of the irregularities $q(V') = \dim H^1(V', \mathcal{O}_{V'})$ for all the finite étale coverings $V' \rightarrow V$.

If X is nonsingular projective with $\kappa(X) = 0$, then $q(X) \leq q^{\max}(X) \leq \dim X$ by [32]. Let V be a normal projective variety with only canonical singularities such that $\kappa(V) = 0$. Let X be a nonsingular projective variety birational to V . Then $q(X) = q(V)$ and $\mathrm{Alb}(X) \simeq \mathrm{Alb}(V)$, since V has only rational singularities. Furthermore, $\pi_1^{\mathrm{alg}}(X) \simeq \pi_1^{\mathrm{alg}}(V)$ by Remark 3.7. In particular, the category of finite étale coverings over X is equivalent to that of finite étale coverings over V . Therefore, $q^{\max}(X) = q^{\max}(V)$.

Definition 4.2. Let F be a normal projective variety with only canonical singularities such that $K_F \sim_{\mathbb{Q}} 0$. If $q^{\max}(F) = 0$, then F is called a *weak Calabi-Yau variety*.

Let V be a normal projective variety with only canonical singularities such that $K_V \sim_{\mathbb{Q}} 0$. Then, by [33, Corollary 8.4], there is a finite étale covering $F \times A \rightarrow V$ with F a weak Calabi-Yau variety and A an abelian variety. Here, $q^{\max}(V) = \dim A$.

The result below guarantees the uniqueness (up to isomorphism) of minimal étale cover $V^{\sim}$ of V realizing $q^{\max}(V)$ as $q(V^{\sim})$. A similar result is found in [4].

Proposition 4.3. *Let V be a normal projective variety with only canonical singularities such that $\kappa(V) = 0$. Then there exists a finite étale Galois covering $V^{\sim} \rightarrow V$, unique up to (non-canonical) isomorphisms, such that:*

- (1) $q(V^{\sim}) = q^{\max}(V)$, and
- (2) *if $V' \rightarrow V$ is a finite étale covering from a variety V' with $q(V') = q^{\max}(V)$, then $V' \rightarrow V$ factors through $V^{\sim} \rightarrow V$.*

Proof. There is a finite étale Galois covering $V_0 \rightarrow V$ with $q^{\max}(V) = q(V_0)$. For the Galois group G_0 of $V_0 \rightarrow V$, let H_0 be the kernel of the natural homomorphism

$$G_0 \rightarrow \mathrm{Aut}(H_1(\mathrm{Alb}(V_0), \mathbb{Z})).$$

We set $V^{\sim}$ to be the quotient space $H_0 \backslash V_0$. Then $H_0 \backslash \mathrm{Alb}(V_0)$ is an abelian variety and $V^{\sim} \rightarrow H_0 \backslash \mathrm{Alb}(V_0)$ is the Albanese map of $V^{\sim}$. In particular, $q^{\max}(V) = q(V^{\sim})$. Let

$V_1 \rightarrow V$ be a finite étale Galois covering such that $V_1 \rightarrow V$ factors as $V_1 \rightarrow V_0 \rightarrow V$. For the Galois group G_1 of $V_1 \rightarrow V$, let $H_1 \subset G_1$ be the kernel of

$$G_1 \rightarrow \text{Aut}(H_1(\text{Alb}(V_1), \mathbb{Z})).$$

Then H_1 is just the pullback of $H_0 \subset G_0$ by $G_1 \rightarrow G_0$, since $\text{Alb}(V_1) \rightarrow \text{Alb}(V_0)$ is an isogeny. In particular, $H_1 \backslash V_1 \simeq H_0 \backslash V_0 = V^\sim$. Therefore, $V^\sim$ is independent of the choice of Galois covering $V_0 \rightarrow V$. For an arbitrary finite étale cover $V' \rightarrow V$ with $q(V') = q^{\max}(V)$, we have a finite Galois cover $V_0 \rightarrow V$ which factors through $V' \rightarrow V$. Then the Galois group G'_0 of $V_0 \rightarrow V'$ acts on $\text{Alb}(V_0)$ and, acts on $H_1(\text{Alb}(V_0), \mathbb{Z})$ trivially, since $\text{Alb}(V_0) \rightarrow \text{Alb}(V')$ is an isogeny. Thus $G'_0 \subset H_0$ and we have a factorization $V' \rightarrow V^\sim \rightarrow V$. $\square$

We call the Galois cover $V^\sim \rightarrow V$ the *Albanese closure*. The Albanese closures of birationally equivalent varieties satisfying the conditions of Proposition 4.3 are also birationally equivalent.

4.2. Splitting endomorphisms. We shall show that any nearly étale rational endomorphism φ of the product $F \times A$ of a weak Calabi-Yau variety F and an abelian variety A is split as the product $\varphi_F \times \varphi_A$ of a nearly étale rational endomorphism φ_F of F and an étale endomorphism φ_A of A . A slightly more general assertion is proved in Proposition 4.8 below. To begin with, we recall the following well-known result (cf. [27, Theorem (4.6)]):

Lemma 4.4. *Let F be a normal projective variety such that $q(F) = 0$. If F is not ruled, then $\text{Aut}(F)$ is discrete.*

Proof. Let $\mathcal{H}$ be a very ample invertible sheaf of F . For an automorphism f of F belonging to the identity component $\text{Aut}^0(F)$, the invertible sheaf $f^*\mathcal{H}$ is isomorphic to $\mathcal{H}$, since the tangent space of the Picard scheme of F at the origin $[\mathcal{O}_F]$ is isomorphic to the zero-dimensional vector space $H^1(F, \mathcal{O}_F)$. Let $\Phi: F \hookrightarrow \mathbb{P}^N$ be the embedding defined by the very ample linear system $|\mathcal{H}|$. Then f induces an automorphism $\rho(f): \mathbb{P}^N \rightarrow \mathbb{P}^N$ such that $\Phi \circ f = \rho(f) \circ \Phi$. The automorphism $\rho(f)$ is contained in a linear subgroup of $\text{PGL}(N+1, \mathbb{C})$ preserving $\Phi(F)$. Since F is not ruled, we infer that the linear subgroup acts on F trivially. Therefore, $f = \text{id}_F$. Hence, $\text{Aut}^0(F) = \{\text{id}_F\}$. $\square$

The following is the first splitting criterion for an étale morphism.

Lemma 4.5. *Let F and F' be non-ruled normal projective varieties such that $q(F) = q(F') = 0$ and $\dim F = \dim F'$. Let A and A' be abelian varieties with $\dim A = \dim A'$. Let $\varphi: F \times A \rightarrow F' \times A'$ be a surjective étale morphism. Then $\varphi = \varphi_1 \times \varphi_2$ for surjective étale morphisms $\varphi_1: F \rightarrow F'$ and $\varphi_2: A \rightarrow A'$.*

Proof. The second projections $p_2: F \times A \rightarrow A$ and $p'_2: F' \times A' \rightarrow A'$ are the Albanese maps of $F \times A$ and $F' \times A'$, respectively. Thus an étale map $\varphi_2: A \rightarrow A'$ is induced so that $\varphi_2 \circ p_2 = p'_2 \circ \varphi$. So for any $a \in A$, there is an étale morphism $\rho_a: F \rightarrow F'$ such that

$$\varphi(x, a) = (\rho_a(x), \varphi_2(a)).$$

The collection $\{\rho_a\}$ gives rise to a morphism from A into the scheme $\text{Hom}(F, F')$ of morphisms from F to F' . For a surjective étale morphism $\psi: F \rightarrow F'$, the tangent space of $\text{Hom}(F, F')$ at the point $[\psi] \in \text{Hom}(F, F')$ corresponding to ψ is isomorphic to

$$\text{Hom}_{\mathcal{O}_F}(\psi^* \Omega_{F'}^1, \mathcal{O}_F) \simeq \text{Hom}_{\mathcal{O}_F}(\Omega_F^1, \mathcal{O}_F) \simeq H^0(F, \Theta_F)$$

for the tangent sheaf Θ_F . In particular, the dimension of the tangent space equals that of $\text{Aut}^0(F)$. So the tangent space is zero by Lemma 4.4. Hence, ρ_a is independent of the choice of $a \in A$. Thus, $\varphi = \varphi_1 \times \varphi_2$ for $\varphi_1 = \rho_a$. $\square$

The following is a partial generalization of Lemma 4.5 in the birational case.

Lemma 4.6. *Let F and F' be non-ruled nonsingular projective varieties such that $q(F) = q(F') = 0$ and $\dim F = \dim F'$. Let A and A' be abelian varieties with $\dim A = \dim A'$. Let $\varphi: F \times A \dashrightarrow F' \times A'$ be a birational map. Then $\varphi = \varphi_1 \times \varphi_2$ for a birational map $\varphi_1: F \dashrightarrow F'$ and an isomorphism $\varphi_2: A \xrightarrow{\sim} A'$.*

Proof. There is an isomorphism $\varphi_2: A \rightarrow A'$ such that $p'_2 \circ \varphi = \varphi_2 \circ p_2$ for the second projections $p_2: F \times A \rightarrow A$ and $p'_2: F' \times A' \rightarrow A'$. Hence, we may assume that $A = A'$ and $\varphi_2 = \text{id}_A$. Then φ is a birational map $F \times A \dashrightarrow F' \times A$ over A . For a general point $a \in A$, we have a birational map $f_a: F \dashrightarrow F'$ as the restriction of φ to $F \times \{a\}$. Thus, we may also replace F' with F by a suitable f_a . Therefore, we may assume from the beginning that φ is a birational map $F \times A \dashrightarrow F \times A$ over A . Then φ induces a rational map from A into the scheme $\text{Bir}(F)$ of birational automorphisms studied in [27]. By [28, Theorem (2.1)], we have $\dim \text{Bir}(F) = 0$, and hence the map $A \dashrightarrow \text{Bir}(F)$ is constant. Therefore, $\varphi = \varphi_F \times \text{id}_A$ for a birational map $\varphi_F: F \dashrightarrow F$. $\square$

The next is a sufficient condition to split the variety into a product.

Lemma 4.7. *Let V be a normal projective variety with only canonical singularities such that $K_V \sim_{\mathbb{Q}} 0$. Suppose that V is birational to $F \times A$ for an abelian variety A and a normal variety F with only canonical singularities such that $K_F \sim_{\mathbb{Q}} 0$ and $q(F) = 0$. Then $V \simeq F' \times A$ for a variety F' birational to F .*

Proof. The composition $V \dashrightarrow F \times A \rightarrow A$ with the second projection is the Albanese map of V . There is a finite Galois étale covering $A' \rightarrow A$ from an abelian variety A' such

that $V \times_A A' \simeq F' \times A'$ over A' for a variety F' , by [33, Theorem 8.3]. Then F' is normal with only canonical singularities, $K_{F'} \sim_{\mathbb{Q}} 0$, $q(F') = 0$, and we have a birational map

$$\varphi: F \times A' = (F \times A) \times_A A' \cdots \rightarrow V \times_A A' \simeq F' \times A'$$

over A' . By Lemma 4.6, $\varphi = \varphi_F \times \text{id}_{A'}$ for a birational map $\varphi_F: F \cdots \rightarrow F'$, since the irregularities of nonsingular models of F and F' are both zero. The action of the Galois group G of $A' \rightarrow A$ on $F' \times A' \simeq V \times_A A'$ is written as a diagonal action by Lemma 4.5. Moreover, it is compatible with the action of G on $F \times A'$ by φ , where G acts trivially on the first factor F . Therefore, G acts trivially on the first factor F' , and hence, $V \simeq F' \times A$. $\square$

The following is also a partial generalization of Lemma 4.5.

Proposition 4.8. *Let F and F' be normal projective varieties with only canonical singularities such that $K_F \sim_{\mathbb{Q}} 0$, $K_{F'} \sim_{\mathbb{Q}} 0$, and $q(F) = q(F') = 0$. Let A and A' be abelian varieties with $\dim A = \dim A'$. Let $\varphi: F \times A \cdots \rightarrow F' \times A'$ be a nearly étale rational map such that $p'_2 \circ \varphi = \varphi_A \circ p_2$ for an étale morphism $\varphi_A: A \rightarrow A'$ and for the second projections $p_2: F \times A \rightarrow A$ and $p'_2: F' \times A' \rightarrow A'$. Then $\varphi = \varphi_F \times \varphi_A$ for a nearly étale rational map $\varphi_F: F \cdots \rightarrow F'$.*

Proof. By Lemma 3.4, φ is the composite of a birational map $F \times A \cdots \rightarrow V^\sharp$ and a finite étale covering $V^\sharp \rightarrow F' \times A'$, since $F' \times A'$ has only canonical singularities (cf. Remark 3.7). Then $V^\sharp \simeq F^\sharp \times A$ for a normal projective variety $F^\sharp$ birational to F by Lemma 4.7. Further, the étale covering $V^\sharp \rightarrow F' \times A'$ is isomorphic to $\psi \times \varphi_A$ for finite étale morphisms $\psi: F^\sharp \rightarrow F'$ and $\varphi_A: A \rightarrow A'$ by Lemma 4.5. So we have only to show that the birational map $F \times A \cdots \rightarrow V^\sharp \simeq F^\sharp \times A$ is the product of a birational map $F \cdots \rightarrow F^\sharp$ and the identity map $A \rightarrow A$. This is done by Lemma 4.6, since the irregularities of nonsingular models of F and $F^\sharp$ are both zero. $\square$

4.3. Proof of Theorem B. As in Proposition 4.8, h is the composite of a birational map $V \cdots \rightarrow V^\sharp$ and an étale covering $\lambda: V^\sharp \rightarrow V$. Let $\delta: V^\sim \rightarrow V$ be the Albanese closure of V and let U be a connected component of the fiber product $V^\sharp \times_V V^\sim$. Then $q(U) = q(V^\sim) = q^{\max}(V)$, and hence $U \rightarrow V^\sharp$ factors through the Albanese closure $(V^\sharp)^\sim \rightarrow V^\sharp$ of $V^\sharp$. On the other hand, $(V^\sharp)^\sim$ is birational to $V^\sim$ by the birational map $V \cdots \rightarrow V^\sharp$. Hence we have $U = V^\sharp \times_V V^\sim \simeq (V^\sharp)^\sim$ by comparing the mapping degrees. So there is a nearly étale rational map $h^\sim: V^\sim \cdots \rightarrow V^\sim$ such that $\delta \circ h^\sim = h \circ \delta$.

Let $\alpha: V^\sim \rightarrow A$ be the Albanese map of $V^\sim$. Then $h^\sim$ induces an étale endomorphism $h_A: A \rightarrow A$ such that $\alpha \circ h^\sim = h_A \circ \alpha$. By [33, Theorem 8.3, Corollary 8.4], there are an isogeny $A' \rightarrow A$ of abelian varieties and an isomorphism $V^\sim \times_A A' \simeq F \times A'$ over

A' for a normal projective variety F with only canonical singularities and $q(F) = 0$. By replacing the isogeny $A' \rightarrow A$, we may assume that $A' = A$ and $A' \rightarrow A$ is the multiplication map by $m > 0$. By Lemma 4.9 below, there is an étale endomorphism φ_A of A such that $h_A(ma) = m\varphi_A(a)$ for any $a \in A$. So there is a nearly étale rational map $\varphi: F \times A \cdots \rightarrow F \times A$ such that $p_2 \circ \varphi = \varphi_A \circ p_2$ and $\theta \circ \varphi = h^\sim \circ \theta$, where $\theta: F \times A \rightarrow V^\sim$ is the composite of the isomorphism $F \times A' \simeq V^\sim \times_A A'$ and the projection $V^\sim \times_A A' \rightarrow V^\sim$. Then $\varphi = \varphi_F \times \varphi_A$ for a nearly étale rational map $\varphi_F: F \cdots \rightarrow F$, by Proposition 4.8. Applying the same argument above to $\varphi_F: F \cdots \rightarrow F$ instead of $h: V \cdots \rightarrow V$, we complete the proof of Theorem B by induction on $\dim V$. $\square$

Lemma 4.9. *Let $\mu: A \rightarrow A$ be the multiplication map $a \mapsto ma$ by a positive integer m for an abelian variety A with a given abelian group structure. Then, for a morphism $h: A \rightarrow A$, there is a morphism $h': A \rightarrow A$ such that $\mu \circ h' = h \circ \mu$.*

Proof. There exist a homomorphism $\varphi: A \rightarrow A$ of abelian group and a point $c \in A$ such that $h(a) = \varphi(a) + c$ for any $a \in A$. There is a point $c' \in A$ with $mc' = c$, since A is divisible. We define $h': A \rightarrow A$ by $h'(a) = \varphi(a) + c'$ for $a \in A$. Then

$$mh'(a) = m\varphi(a) + mc' = \varphi(ma) + c = h(ma). \quad \square$$

4.4. Conjectural discussion. We shall pose the following:

Conjecture 4.10. Let X be a nonsingular projective variety such that $\kappa(X) = q^{\max}(X) = 0$. Then $\pi_1(X)$ is finite.

This is true for X with K_X numerically trivial, by Bogomolov's decomposition theorem. Furthermore, it is true when $\dim X \leq 3$ (cf. [50]).

Lemma 4.11. *Let V be a normal projective variety such that $\pi_1^{\text{alg}}(M)$ is finite for a resolution $M \rightarrow V$ of singularities of V . Then any nearly étale rational endomorphism $h: V \cdots \rightarrow V$ is birational.*

Proof. We may assume that V is nonsingular. For an arbitrary positive integer l , let $V \cdots \rightarrow V^{\sharp(l)} \rightarrow V$ be the Stein factorization of the l -th power $h^l = h \circ \cdots \circ h$. Then $V^{\sharp(l)} \rightarrow V$ is an étale morphism of degree $(\deg h)^l$ by Lemma 3.4. Now, $\pi_1^{\text{alg}}(V)$ is finite, since $\pi_1(M) \simeq \pi_1(V)$. Hence $\deg h = 1$. $\square$

So, if Conjecture 4.10 is true, then a nearly étale rational endomorphism of a weak Calabi-Yau variety is birational. In particular, the *building blocks* of the étale endomorphisms of projective varieties with $\kappa = 0$ would then turn out to be the endomorphisms of abelian varieties and the birational automorphisms of weak Calabi-Yau varieties.

5. THE UNIRULED CASE

5.1. Maximal rationally connected fibration. A projective variety X is called *uniruled* if there is a dominant rational map $\mathbb{P}^1 \times Y \dashrightarrow X$ for a variety Y with $\dim Y = \dim X - 1$. For a nonsingular projective variety X , the following three conditions are all equivalent (cf. [9, §3], [40, §2]):

- (1) Any two points of X are connected by a chain of rational curves.
- (2) Any two general points of X are contained in one and the same rational curve.
- (3) There is a nonsingular rational curve on X with ample normal bundle.

If one of the conditions above is satisfied, then X is called *rationally connected*.

Remark 5.1 (cf. [9, §3], [40, Proposition (2.5)]). A nonsingular rationally connected variety X has the following properties:

- X is simply connected.
- $H^i(X, \mathcal{O}_X) = 0$ for $i > 0$.
- $H^0(X, (\Omega_X^1)^{\otimes m}) = 0$ for any $m > 0$.

If X is a uniruled nonsingular projective variety, then there exists uniquely up to birational equivalence, a rational fiber space $\pi: X \dashrightarrow Y$ over a nonsingular projective variety Y satisfying the following conditions, by [10], [40], [22]:

- (1) π is weakly holomorphic, i.e., there exists open dense subsets $U \subset X$, $V \subset Y$ such that π induces a proper surjective morphism $U \rightarrow V$.
- (2) For a general point P of the open set U above, the fiber over $\pi(P)$ is a maximal rationally connected submanifold of X containing P .
- (3) Y is not uniruled (cf. [22, Corollary 1.4]).

The fibration π is called the *maximal rationally connected fibration* of X .

Lemma 5.2. *Let $f: X \rightarrow X$ be an étale endomorphism of a uniruled projective variety X . Then there exist a proper birational morphism $\mu: M \rightarrow X$, a proper surjective morphism $\pi: M \rightarrow Y$, an étale endomorphism f_M of M , and an endomorphism h of Y satisfying the following conditions:*

- (1) M is a nonsingular projective variety.
- (2) Y is a normal and non-uniruled projective variety.
- (3) π is birational to the maximal rationally connected fibration of M .
- (4) $\mu \circ f_M = f \circ \mu$ and $\pi \circ f_M = h \circ \pi$ (cf. the diagram in Theorem C).

Proof. We may assume that X is nonsingular, by replacing it with an equivariant resolution of singularities with respect to f (cf. Section 1.4). Let $\Psi: X \dashrightarrow \text{Chow}(X)$ be the rational map to the Chow variety of X which defines the maximal rationally connected

fibration of X . By associating a cycle Z of X with the push-forward f_*Z , we have a functorial morphism $f_c: \text{Chow}(X) \rightarrow \text{Chow}(X)$ since f is finite (cf. [3, Chapter IV §2 Théorème 6], [2, Théorème 6.3.1], [38, Theorem I.6.8]). Since f is étale, f_*Z is reduced for a rationally connected submanifold Z . Thus, $\Psi \circ f = f_c \circ \Psi$. Let $X \cdots \rightarrow Y \rightarrow \text{Chow}(X)$ be the Stein factorization of Ψ and $h: Y \rightarrow Y$ the endomorphism induced from f_c and f . Note that Y is not uniruled. An endomorphism of the graph $\Gamma \subset X \times Y$ of the rational map $X \cdots \rightarrow Y$ is induced from $f \times h$. Let $T \rightarrow \Gamma$ be the normalization, f_T the induced endomorphism of T , and $p_1: T \rightarrow X$ the morphism induced from the first projection $\Gamma \subset X \times T \rightarrow T$. Then, $p_1 \circ f_T = f \circ p_1$. Thus, p_1 factors through a finite birational morphism $T \rightarrow T_1$ to the fiber product $T_1 := X \times_X T$ of f and p_1 over X . Since f is étale, $T \simeq T_1$, and hence f_T is étale. Let $M \rightarrow T$ be an equivariant resolution of singularities with respect to f_T , and f_M the lift of f_T to M as an étale endomorphism. Then, these data satisfy the required conditions. $\square$

The following is proved essentially in [37, Theorem 5.2], where F is assumed to be a rationally connected manifold. We present a slightly different proof.

Lemma 5.3. *Let $g: M \rightarrow N$ be a proper surjective morphism between nonsingular varieties. For a general fiber F of g , suppose that*

- (1) F is connected,
- (2) F is simply connected, and
- (3) $H^i(F, \mathcal{O}_F) = 0$ for any $i > 0$.

Then $g_*: \pi_1(M) \rightarrow \pi_1(N)$ is isomorphic.

Proof. Step 1. If $N^\circ \subset N$ is a Zariski open subset with the codimension of $N \setminus N^\circ$ bigger than one, then $\pi_1(N^\circ) \simeq \pi_1(N)$, and $\pi_1(g^{-1}(N^\circ)) \rightarrow \pi_1(M)$ is surjective. Thus, if $\pi_1(g^{-1}(N^\circ)) \rightarrow \pi_1(N^\circ)$ is an isomorphism, then so is $\pi_1(M) \rightarrow \pi_1(N)$. Hence, we may replace N with such an open subset N° . In particular, we may assume that $g: M \rightarrow N$ is smooth outside a nonsingular divisor $D = \sum D_i$ of N , where D_i is an irreducible component.

Step 2. Let $\mathcal{U}_i$ be an analytic open neighborhood of a point $P_i \in D_i$ of D such that $\mathcal{U}_i$ is biholomorphic to a unit polydisc $\{(t_1, t_2, \dots, t_n) \in \mathbb{C}^n; |t_j| < 1 \text{ for any } j\}$ in which P_i is mapped to the origin $0 = (0, 0, \dots, 0)$ and $\mathcal{U}_i \cap D$ is mapped to the coordinate hypersurface $\{t_1 = 0\}$. Since $\mathcal{U}_i \setminus D = \mathcal{U}_i \setminus D_i$ is homotopic to a circle, we have a generator δ_i of $\pi_1(\mathcal{U}_i \setminus D)$. By van Kampen's theorem, or other topological argument, we infer that the kernel of the surjection $\pi_1(N \setminus D) \rightarrow \pi_1(N)$ is generated by the conjugacy classes of the images $\bar{\delta}_i$ of δ_i under the homomorphisms $\pi_1(\mathcal{U}_i \setminus D) \rightarrow \pi_1(N \setminus D)$.

Step 3. By the assumptions (1) and (2), and by the homotopy exact sequence, we infer that the natural homomorphism $\pi_1(M \setminus g^{-1}D) \rightarrow \pi_1(N \setminus D)$ is an isomorphism. Let $\hat{\delta}_i \in \pi_1(M \setminus g^{-1}D)$ be the element corresponding to $\bar{\delta}_i \in \pi_1(N \setminus D)$. In order to show the homomorphism $\pi_1(M) \rightarrow \pi_1(N)$ to be isomorphic, it is enough to show that $\hat{\delta}_i$ is contained in the kernel of $\pi_1(M \setminus g^{-1}D) \rightarrow \pi_1(M)$. Let $C_i \subset \mathcal{U}_i$ be a curve corresponding to an axis with respect to the coordinate system $(t_1, \dots, t_n)$ of the polydisc such that $C_i \cap D = \{P_i\}$ and let X_i be the fiber product $M \times_N C_i$. By changing P_i , $\mathcal{U}_i$, and coordinates $(t_1, \dots, t_n)$ slightly, we may assume that X_i is nonsingular. Then $\hat{\delta}_i$ comes from $\pi_1(X_i \setminus g^{-1}(P_i)) \simeq \pi_1(C_i \setminus \{P_i\}) = \mathbb{Z}\delta_i$. Thus, we have only to show that $\pi_1(X_i) = 0$, or equivalently, $\pi_1(g^{-1}(P_i)) = 0$, since $g^{-1}(P_i)$ is a deformation retract of X_i .

Step 4. By *Step 3*, the proof of Lemma 5.3 is reduced to the case where N is a unit disc and g is smooth outside the origin 0. We shall show $\pi_1(g^{-1}(0)) = 0$ in this case. By shrinking N if necessary, we have a holomorphic curve $T \subset M$ such that T is biholomorphic to a unit disc and $T \rightarrow N$ is a finite surjective morphism branched only at $0 \in N$. We have:

- $\pi_1(M \setminus g^{-1}(0)) \simeq \pi_1(N \setminus \{0\}) \simeq \mathbb{Z}$,
- $\pi_1(T \setminus g^{-1}(0)) \rightarrow \pi_1(N \setminus \{0\})$ is an injection into a finite-index subgroup, and
- $\pi_1(M \setminus g^{-1}(0)) \rightarrow \pi_1(M)$ is surjective.

Therefore, $\pi_1(M)$ is a finite cyclic group. Let $\lambda: \widetilde{M} \rightarrow M$ be the universal covering map and let $\widetilde{M} \rightarrow \widetilde{N} \rightarrow N$ be the Stein factorization of $g \circ \lambda: \widetilde{M} \rightarrow M \rightarrow N$. Then the induced morphism $\widetilde{M} \rightarrow M \times_N \widetilde{N}$ is isomorphic over $(N \setminus \{0\}) \times_N \widetilde{N}$ by (2). Applying Lemma 3.9 to the commutative diagram corresponding to the Stein factorization of $g \circ \lambda$, we infer that $\widetilde{N} \rightarrow N$ is étale. In fact, the condition (5) of Lemma 3.9 over N and $\widetilde{N}$ are both satisfied, since the higher direct image sheaves $R^i g_* \mathcal{O}_M$ are torsion free over the nonsingular curves N and the same thing holds for the higher direct image sheaves for $\widetilde{M} \rightarrow \widetilde{N}$. Hence, $\widetilde{N} \simeq N$ and $\widetilde{M} \simeq M$. Therefore, M and $g^{-1}(0)$ are simply connected. $\square$

The following gives a sufficient condition for a finite morphism to be nearly étale.

Proposition 5.4. *Let $h: \widetilde{Y} \rightarrow Y$ be a finite surjective morphism between normal varieties with $\deg h > 1$. Then h is nearly étale if there exist proper surjective morphisms $\pi: M \rightarrow Y$ and $\tilde{\pi}: \widetilde{M} \rightarrow \widetilde{Y}$, and a finite étale covering $f: \widetilde{M} \rightarrow M$ satisfying the following conditions:*

- (1) *M and $\widetilde{M}$ are nonsingular, and $\pi \circ f = h \circ \tilde{\pi}$.*
- (2) *A general fiber F of π is connected and simply connected with $H^i(F, \mathcal{O}_F) = 0$ for any $i > 0$.*

Remark 5.5. Let M_1 be the fiber product $\tilde{Y} \times_Y M$. Then f induces a finite morphism $\lambda: \tilde{M} \rightarrow M_1$ over M with the commutative diagram below:

$$\begin{array}{ccccc} \tilde{M} & \xrightarrow{\lambda} & M_1 & \longrightarrow & M \\ \tilde{\pi} \downarrow & & \downarrow & & \downarrow \pi \\ \tilde{Y} & \xlongequal{\quad} & \tilde{Y} & \xrightarrow{h} & Y. \end{array}$$

We can show that λ is just the normalization of the reduced structure $M_{1,\text{red}}$ of M_1 , as follows: Note that M_1 is irreducible since π has only connected fibers and h is finite. Since a general fiber of π is simply connected, λ is an isomorphism over $h^{-1}(U) \times_Y M \subset \tilde{Y} \times_Y M = M_1$ for a dense open subset $U \subset Y$. Then λ induces a finite and birational morphism from $\tilde{M}$ to the normalization of $M_{1,\text{red}}$, which is turned to be an isomorphism.

Proof of Proposition 5.4. Let $\nu: N \rightarrow Y$ be a resolution of singularities and $M' \rightarrow M$ a proper birational morphism from a nonsingular variety M' such that $M' \rightarrow M \cdots \rightarrow N$ is a morphism. Then, $\pi_1(M') \simeq \pi_1(N)$ by Lemma 5.3. Let $\Pi \subset \pi_1(N)$ be the image of $f_*(\pi_1(\tilde{M})) \subset \pi_1(M)$ via $\pi_1(M) \simeq \pi_1(M') \simeq \pi_1(N)$, and $\phi: \tilde{N} \rightarrow N$ the finite étale covering corresponding to the finite-index subgroup Π . Let $\tilde{M}' \rightarrow M'$ be the étale covering obtained as the pullback of f by $M' \rightarrow M$. Then we have a commutative diagram

$$\begin{array}{ccccc} \tilde{M} & \longleftarrow & \tilde{M}' & \longrightarrow & \tilde{N} \\ f \downarrow & & \downarrow & & \downarrow \phi \\ M & \longleftarrow & M' & \longrightarrow & N \end{array}$$

in which each square is Cartesian. By considering the Stein factorization of $\nu \circ \phi$, we have a proper birational morphism $\tilde{\nu}: \tilde{N} \rightarrow \tilde{Y}$ such that $h \circ \tilde{\nu} = \nu \circ \phi$. Thus, the diagram

$$\begin{array}{ccccc} \tilde{M} & \xrightarrow{\tilde{\pi}} & \tilde{Y} & \xleftarrow{\tilde{\nu}} & \tilde{N} \\ f \downarrow & & h \downarrow & & \downarrow \phi \\ M & \xrightarrow{\pi} & Y & \xleftarrow{\nu} & N \end{array}$$

is also commutative. Therefore, h is nearly étale. $\square$

5.2. Proof of Theorem C. We apply Lemma 5.2 to the given étale endomorphism f of X . Let $\mu: M \rightarrow X$, $\pi: M \rightarrow Y$, f_M , and h be the same objects as in Lemma 5.2. By Proposition 5.4, h is nearly étale. This completes the proof of Theorem C. $\square$

APPENDIX A. TOPOLOGICAL ENTROPIES AND FIBER SPACES

The reduction (A) in the introduction is deduced from Theorem A. We have also a similar reduction of type (A) in the dynamical study of endomorphisms. More precisely,

for the study of the first dynamical degree and the topological entropies of holomorphic surjective endomorphisms of compact Kähler manifolds, the case of positive Kodaira dimension is reduced to the case of zero Kodaira dimension by Theorem A, and by Theorem D below. The reduction of non-holomorphic meromorphic endomorphisms is not treated here.

Theorem D. *Let $\pi: X \rightarrow Y$ be a proper surjective morphism from a compact Kähler manifold X to a compact complex analytic variety Y and $f: X \rightarrow X$ a surjective endomorphism such that a general fiber of π is connected and $\pi \circ f = \pi$. Let F be a smooth fiber of π , i.e., a fiber along which π is smooth. Then:*

- (1) *The equality $d_1(f) = d_1(f|_F)$ holds for the first dynamical degrees d_1 of $f: X \rightarrow X$ and the restriction $f|_F: F \rightarrow F$ of f .*
- (2) *If f is étale, then the equality $h_{\text{top}}(f) = h_{\text{top}}(f|_F)$ holds for the topological entropy h_{top} .*

The proof is based on basic properties of spectral radii, a generalized notion of Kähler cone, a simple calculation of dynamical degrees, and the results of Gromov [24] and Yomdin [56] on topological entropies. The proof of the first assertion (1) is given at the end of Section A.2 and the proof of the second assertion (2) at the end of Section A.3. Some well-known properties on dynamical degrees are prepared in Sections A.1 and A.2.

A.1. Spectral radii and Kähler cones. We recall some basic properties of spectral radii, especially a generalization of the Perron-Frobenius theorem. Furthermore, we introduce a notion of Kähler (k, k) -forms and the Kähler cones in $H^{k,k}(M, \mathbb{R})$ for a compact Kähler manifold M .

The spectral radius $\rho(\varphi) = \rho(V, \varphi)$ of an endomorphism $\varphi: V \rightarrow V$ of a finite-dimensional $\mathbb{C}$ -vector space V is defined to be the maximum of the absolute values of the eigenvalues of φ . The spectral radius of an endomorphism $\varphi_{\mathbb{R}}: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ of a finite-dimensional real vector space $V_{\mathbb{R}}$ is defined as that of its complexification $\varphi_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}: V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C} \rightarrow V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$.

Remark. Let $\|\cdot\|$ be any norm of V and let $\|\cdot\|_1$ be the L^1 -norm of $\text{End}_{\mathbb{C}}(V)$ defined by $\|\varphi\|_1 := \sup\{\|\varphi(v)\|; \|v\| = 1\}$. Then $\rho(\varphi) = \lim_{m \rightarrow \infty} (\|\varphi^m\|_1)^{1/m}$.

Notation. Let $V_{\mathbb{R}}$ be a finite-dimensional real vector space. A subset $\mathcal{C} \subset V_{\mathbb{R}}$ is called a *convex cone* if $\mathcal{C} + \mathcal{C} \subset \mathcal{C}$ and $\mathbb{R}_+ \mathcal{C} \subset \mathcal{C}$, where $\mathbb{R}_+$ denotes the set of positive real numbers. If $\mathcal{C} \cap (-\mathcal{C}) \subset \{0\}$ in addition, then $\mathcal{C}$ is called *strictly convex*.

Remark. A convex cone $\mathcal{C} \subset V_{\mathbb{R}}$ is strictly convex if and only if there exists a linear form $\chi: V_{\mathbb{R}} \rightarrow \mathbb{R}$ such that $\chi > 0$ on $\mathcal{C} \setminus \{0\}$.

The following is known as a generalization of the Perron-Frobenius theorem on real $n \times n$ matrices $A = (a_{ij})$ of positive entries a_{ij} :

Theorem A.1 (cf. [7]). *Let $\mathcal{C}$ be a strictly convex closed cone of a finite-dimensional real vector space $V_{\mathbb{R}}$ such that $\mathcal{C}$ generates $V_{\mathbb{R}}$ as a vector space. Let $\varphi: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ be an endomorphism such that $\varphi(\mathcal{C}) \subset \mathcal{C}$. Then the spectral radius $\rho(\varphi)$ is an eigenvalue of φ and there is an eigenvector in $\mathcal{C}$ with the eigenvalue $\rho(\varphi)$.*

We add another property of spectral radius which is proved by using Theorem A.1:

Proposition A.2. *Let $\varphi: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ be an endomorphism of a finite-dimensional real vector space $V_{\mathbb{R}}$. Suppose that $\varphi(\mathcal{C}) \subset \mathcal{C}$ for a strictly convex closed cone $\mathcal{C} \subset V_{\mathbb{R}}$ with $\mathcal{C} + (-\mathcal{C}) = V_{\mathbb{R}}$. Let $\|\cdot\|$ be a norm on $V_{\mathbb{R}}$ and let $\chi: V_{\mathbb{R}} \rightarrow \mathbb{R}$ be a linear form such that $\chi > 0$ on $\mathcal{C} \setminus \{0\}$. If u is contained in the interior of $\mathcal{C}$, then*

$$\rho(\varphi) = \lim_{m \rightarrow \infty} \|\varphi^m(u)\|^{1/m} = \lim_{m \rightarrow \infty} \chi(\varphi^m(u))^{1/m}.$$

Proof. We define $\|\cdot\|_u: V_{\mathbb{R}} \rightarrow \mathbb{R}_{\geq 0}$ by $\|x\|_u := \inf\{r \geq 0 \mid x + ru \in \mathcal{C}, -x + ru \in \mathcal{C}\}$. Then $\|\cdot\|_u$ is a norm of $V_{\mathbb{R}}$. There exist positive constants C_1, C_2 such that $C_1\|x\| \leq \|x\|_u \leq C_2\|x\|$ for any $x \in V$. Hence, we may assume that $\|\cdot\| = \|\cdot\|_u$. For the induced L^1 -norm $\|\cdot\|_{u,1}$, we have $\|\varphi^m\|_{u,1} \geq \|\varphi^m(u)\|_u$ for $m \geq 1$, since $\|u\|_u = 1$. Applying χ to $\|\varphi^m(u)\|_u u - \varphi^m(u) \in \mathcal{C}$, we have $\|\varphi^m(u)\|_u \chi(u) \geq \chi(\varphi^m(u))$. We have an eigenvector $v \in \mathcal{C}$ with the eigenvalue $\rho(\varphi)$ by Theorem A.1. Applying $\chi \circ \varphi^m$ to $\|v\|_u u - v \in \mathcal{C}$, we have $\|v\|_u \chi(\varphi^m(u)) \geq \rho(\varphi)^m \chi(v)$. Therefore,

$$\|\varphi^m\|_{u,1} \geq \|\varphi^m(u)\|_u \geq \frac{\chi(\varphi^m(u))}{\chi(u)} \geq \frac{\chi(v)}{\chi(u)\|v\|_u} \rho(\varphi)^m.$$

Hence, we complete the proof by

$$\rho(\varphi) = \lim_{m \rightarrow \infty} \|\varphi^m\|_{u,1}^{1/m} \geq \lim_{m \rightarrow \infty} \|\varphi^m(u)\|_u^{1/m} \geq \lim_{m \rightarrow \infty} \chi(\varphi^m(u))^{1/m} \geq \rho(\varphi). \quad \square$$

Let M be a compact Kähler manifold of dimension n . Let ω be a C^∞ -(k, k)-form on M for an integer $1 \leq k \leq n$. For a local coordinate system $(z_1, z_2, \dots, z_n)$ of M , ω is locally expressed as

$$\omega = (\sqrt{-1})^{k^2} \sum_{I, J \subset \{1, 2, \dots, n\}} a_{I, J} dz_I \wedge d\bar{z}_J$$

for C^∞ -functions $a_{I, J}$, where $\sharp I = \sharp J = k$ and

$$dz_I := dz_{i_1} \wedge dz_{i_2} \wedge \cdots \wedge dz_{i_k}$$

when $I = \{i_1, i_2, \dots, i_k\}$ with $i_1 < i_2 < \cdots < i_k$. The ω is called a *Kähler* (k, k)-form if ω is d -closed ($d\omega = 0$) and real ($\bar{\omega} = \omega$), and if the matrix $(a_{I, J})$ is positive-definite everywhere in M . Note that this is just the usual definition of Kähler forms in case $k = 1$.

The following is easily shown:

Lemma A.3.

- (1) If η is a (usual) Kähler form $((1,1)$ -form), then $\eta^k = \eta \wedge \cdots \wedge \eta$ is a Kähler (k,k) -form for $1 \leq k \leq n$.
- (2) Let ω be a Kähler (k,k) -form. If ω' is a Kähler $(n-k, n-k)$ -form, then $\int_M \omega \wedge \omega' > 0$. If T is a d -closed positive $(n-k, n-k)$ -current, then $\int_M \omega \wedge T \geq 0$.

Inside of the real vector space $H^{k,k}(M, \mathbb{R}) := H^{k,k}(M) \cap H^{2k}(M, \mathbb{R})$, the set $P^k(M)$ of the classes $[\omega]$ of Kähler (k,k) -forms ω on M is a strictly convex open cone. It is called the *Kähler cone* of degree k . Its closure in $H^{k,k}(M, \mathbb{R})$ is denoted by $\overline{P^k(M)}$.

- Lemma A.4.** (1) $\int_M \xi \cup [\omega] > 0$ for any $\xi \in \overline{P^k(M)} \setminus \{0\}$ and for any Kähler $(n-k, n-k)$ -form ω .
- (2) If $\theta \in \overline{P^1(M)}$ and if $\theta - [T] \in P^1(M)$ for a d -closed positive $(1,1)$ -current T , then for any $1 \leq l \leq n$, there exists an element $z \in \overline{P^{l-1}(M)}$ such that $\theta^l - z \cup [T] \in P^l(M)$ and $z \cup [T]$ is represented by a d -closed positive (l,l) -current. In particular, $\int_M \xi \cup \theta^{n-k} > 0$ for any $\xi \in \overline{P^k(M)} \setminus \{0\}$.

Proof. (1): Let (x, y) denote $\int_M x \cup y$ for $x \in H^{k,k}(M, \mathbb{R})$ and $y \in H^{n-k, n-k}(M, \mathbb{R})$. Then $H^{n-k, n-k}(M, \mathbb{R})$ is dual to $H^{k,k}(M, \mathbb{R})$ by $(*, *)$. Since $P^{n-k}(M)$ generates $H^{n-k, n-k}(M, \mathbb{R})$ as the vector space, we can find a Kähler $(n-k, n-k)$ -form ω_0 such that $(\xi, [\omega_0]) \neq 0$. By Lemma A.3, we have $(\xi, [\omega_0]) > 0$. There is a positive constant C such that $C\omega - \omega_0$ is also a Kähler (k,k) -form, since M is compact. Thus, by Lemma A.3,

$$(\xi, [\omega]) \geq C^{-1}(\xi, [\omega_0]) > 0.$$

(2): We set $\alpha = \theta - [T] \in P^1(M)$. Then, we have $\theta^l = \alpha^l + z \cup [T]$, where $z = 1 \in H^0(M, \mathbb{R})$ when $l = 1$ and

$$(A-1) \quad z = \sum_{i=0}^{l-1} \theta^i \cup \alpha^{l-1-i}$$

when $l \geq 2$. Thus, $z \in \overline{P^{l-1}(M)}$ and $z \cup [T]$ is represented by a positive current. In particular, $(\xi, \theta^{n-k}) \geq (\xi, \alpha^{n-k}) > 0$ by Lemma A.3 and (1). $\square$

The following is a property of Kähler $(1,1)$ -forms.

Lemma A.5. Let $\psi: M \rightarrow X$ be a generically finite morphism into another compact Kähler manifold X . If η is a Kähler form on X , then $[\psi^*\eta] - [T] \in P^1(M)$ for a d -closed real positive $(1,1)$ -current T on M .

Proof. Let $M \rightarrow V \rightarrow X$ be the Stein factorization of ψ . By Hironaka's blowing up, we have a projective bimeromorphic morphism $\nu: Z \rightarrow V$ from another compact Kähler manifold Z such that $\mu: Z \rightarrow V \cdots \rightarrow M$ is holomorphic. Here, $\mathcal{O}_Z(-E)$ is ν -ample for

a ν -exceptional effective divisor E . Then $\mathcal{O}_Z(-E)$ is also relatively ample over X , and $[\mu^*\psi^*\eta] - \varepsilon[E]$ is represented by a Kähler form on Z for some $\varepsilon > 0$ (cf. [18, Lemma 4.4], [17, Lemma 2]). Thus

$$[\mu^*\psi^*\eta] - \varepsilon[E] - [\mu^*\xi] \in P^1(Z)$$

for a Kähler form ξ on M . Hence, $\psi^*[\eta] - [\xi]$ is represented by a d -closed real positive $(1, 1)$ -current on M . $\square$

A.2. Dynamical degrees. Let $f: M \rightarrow M$ be a surjective endomorphism of a compact Kähler manifold M of dimension n . Then f induces natural homomorphisms

$$f^*: H^i(M, \mathbb{Z}) \rightarrow H^i(M, \mathbb{Z}) \quad \text{and} \quad f_*: H_i(M, \mathbb{Z}) \rightarrow H_i(M, \mathbb{Z})$$

for $0 \leq i \leq 2n$. The composite

$$H^i(M, \mathbb{Z}) \xrightarrow{f^*} H^i(M, \mathbb{Z}) \simeq H_{2n-i}(M, \mathbb{Z}) \xrightarrow{f_*} H_{2n-i}(M, \mathbb{Z}) \simeq H^i(M, \mathbb{Z})$$

is just the multiplication map by $\deg f$, where the isomorphism above is induced from the Poincaré duality. Thus $f^*: H^i(M, \mathbb{R}) \rightarrow H^i(M, \mathbb{R})$ is isomorphic. Moreover, f^* preserves the Hodge structure, i.e., $f^* H^{p,q}(M) = H^{p,q}(M)$. Note that f is a finite morphism, since $f^*: H^{1,1}(M, \mathbb{R}) \rightarrow H^{1,1}(M, \mathbb{R})$ is an isomorphism.

Definition A.6 (Dynamical degree). Let η be a Kähler form on M . For an integer $1 \leq l \leq n$, we set

$$\delta_l(f, \eta) := \int_M f^* \eta^l \wedge \eta^{n-l},$$

where η^i denotes the i -th power $\eta \wedge \cdots \wedge \eta$ for $1 \leq i \leq n$, and $\eta^0 := 1$. The l -th *dynamical degree* of f is defined to be

$$d_l(f) := \lim_{m \rightarrow \infty} (\delta_l(f^m, \eta))^{1/m}.$$

Note that $d_0(f) = 1$ and $d_n(f) = \deg f$ by the definition.

Lemma A.7. Let x be an element of $\overline{P^1(M)}$ and y an element of $\overline{P^k(M)}$ such that $x - [T_1] \in P^1(M)$ and $y - [T_k] \in P^k(M)$ for certain d -closed real positive currents T_1 of type $(1, 1)$ and T_k of type (k, k) for $1 \leq k \leq n$. Then

$$d_{n-k}(f) = \lim_{m \rightarrow \infty} \left(\int_M (f^m)^*(x^{n-k}) \cup y \right)^{1/m}.$$

Proof. There exist d -closed positive currents S_1 of type $(1, 1)$ and S_k of type (k, k) , and a constant $a > 0$ such that $ax - [\eta] = [S_1]$ and $ay - [\eta^k] = [S_k]$. We set $l := n - k$. Then $a^l x^l - [\eta]^l = z \cup [S_1]$ for an element $z \in \overline{P^{l-1}(M)}$ by Lemma A.4, (2). Thus $f^*(a^l x^l - [\eta]^l) = f^*z \cup f^*[S_1]$ is represented by a d -closed positive (l, l) -current, since $f^*[S_1] = [f^*S_1]$ for the positive $(1, 1)$ -current f^*S_1 . Therefore,

$$f^*(a^l x^l) \cup (ay) - f^*[\eta]^l \cup [\eta]^k = f^*z \cup f^*[S_1] \cup (ay) + f^*[\eta]^l \cup [S_k]$$

is represented by a positive (n, n) -current. Hence, for any $m \geq 1$,

$$(A-2) \quad a^{l+1} \int_M (f^m)^*(x^l) \cup y \geq \int_M (f^m)^* \eta^l \wedge \eta^k = \delta_l(f^m, \eta).$$

Conversely, there exists a constant $b > 0$ such that $b[\eta] - x \in P^1(M)$ and $b[\eta^k] - y \in P^k(M)$. Then $b^l[\eta]^l - x^l \in \overline{P^l(M)}$ (cf. (A-1)) and hence $f^*(b^l[\eta]^l - x^l) \in \overline{P^l(M)}$. Therefore

$$f^*(b^{l+1}[\eta]^l) \cup [\eta]^k - f^*(x^l) \cup y \in \overline{P^n(M)}.$$

Thus, for any $m \geq 1$, we have

$$(A-3) \quad b^{l+1} \delta_l(f^m, \eta) = b^{l+1} \int_M (f^m)^* \eta^l \wedge \eta^k \geq \int_M (f^m)^*(x^l) \cup y.$$

Hence, we have the expected equality by applying $m \rightarrow \infty$ to the m -th roots of the both sides of the inequalities (A-2) and (A-3) $\square$

Lemma A.8. *Suppose that there exist a generically finite surjective morphism $\mu: M \rightarrow X$ into another compact Kähler manifold X and an endomorphism $g: X \rightarrow X$ satisfying $\mu \circ f = g \circ \mu$. Then $d_l(f) = d_l(g)$ for any l .*

Proof. Let ξ be a Kähler form on X . By Lemma A.5, $[a\mu^*\xi - \eta]$ is represented by a d -closed real positive $(1, 1)$ -current on M for a certain positive constant a . Then $[a^l\mu^*\xi^l - \eta^l]$ is represented by a d -closed real positive (l, l) -current by Lemma A.4, (2). Thus, by Lemma A.7,

$$\begin{aligned} d_l(f) &= \lim_{m \rightarrow \infty} \left(\int_M (f^m)^*(\mu^*\xi^l) \wedge \mu^*\xi^{n-l} \right)^{1/m} \\ &= \lim_{m \rightarrow \infty} \left((\deg \mu) \int_X (g^m)^*\xi^l \wedge \xi^{n-k} \right)^{1/m} = d_l(g). \end{aligned} \quad \square$$

Lemma A.9 (cf. [13], [26]). *The k -th dynamical degree $d_k(f)$ equals the spectral radius of the following endomorphism induced from f^* :*

$$(f^*)^{(k,k)}: H^{k,k}(M, \mathbb{R}) \rightarrow H^{k,k}(M, \mathbb{R}).$$

Proof. For a Kähler form η , $[\eta^k] \in H^{k,k}(M, \mathbb{R})$ is contained in $P^k(M)$. We define a linear form $\chi: H^{k,k}(M, \mathbb{R}) \rightarrow \mathbb{R}$ by $\chi(x) = \int_M x \cup [\eta^{n-k}]$. Then $\delta_k(f, \eta) = \chi([f^*\eta^k])$. We have $\chi > 0$ on $\overline{P^k(M)} \setminus \{0\}$ by Lemma A.4. Moreover, $(f^*)^{(k,k)}P^k(M) \subset \overline{P^k(M)}$. Therefore, by Proposition A.2,

$$\rho((f^*)^{(k,k)}) = \lim_{m \rightarrow \infty} \chi([(f^m)^*\eta^k])^{1/m} = \lim_{m \rightarrow \infty} \delta_k(f^m, \eta)^{1/m} = d_k(f). \quad \square$$

Fact A.10. We have $d_{l-1}(f)d_{l+1}(f) \leq d_l(f)^2$ for any $1 \leq l < n$ (cf. [25, Proposition 1.2]). In particular, $d_n(f) = \deg f \leq d_1(f)^n$.

For an element $x \in H^{k,k}(M, \mathbb{R})$ and for $l \leq n - k$, we set

$$\delta_l(f, \eta; x) := \int_M [f^* \eta^l \wedge \eta^{n-k-l}] \cup x.$$

Then $\delta_l(f, \eta; [\eta^k]) = \delta_l(f, \eta)$. Let $C > 0$ be a constant such that $C[\eta^k] - x \in P^k(M)$. Then, $C\delta_l(f, \eta) \geq \delta_l(f, \eta; x)$ for any $f: M \rightarrow M$. Hence, we have

$$(A-4) \quad d_l(f) \geq \overline{\lim}_{m \rightarrow \infty} (\delta_l(f^m, \eta; x))^{1/m}.$$

Proposition A.11. *Let F be a compact Kähler manifold of dimension k and let $\phi: F \rightarrow M$ be a generically finite morphism such that $\phi \circ h = f \circ \phi$ for a surjective endomorphism $h: F \rightarrow F$. Then for any $1 \leq l \leq k$, one has*

$$d_l(f) \geq d_l(h) \quad \text{and} \quad d_{l+n-k}(f) \geq \deg(f) \deg(h)^{-1} d_l(h).$$

Proof. Let $G_i: H^{2i}(F, \mathbb{C}) \rightarrow H^{2(n-k+i)}(M, \mathbb{C})$ be the Gysin homomorphism

$$H^{2i}(F, \mathbb{C}) \simeq H_{2k-2i}(F, \mathbb{C}) \xrightarrow{\phi_*} H_{2k-2i}(M, \mathbb{C}) \simeq H^{2n-2k+2i}(M, \mathbb{C})$$

for $0 \leq i \leq k$, where the isomorphisms in both sides are induced from the Poincaré duality. Note that $G_i H^{p,q}(F) \subset H^{p+n-k, q+n-k}(M)$. We have the “projection formula” $G_{p+q}(x \cup \phi^*(y)) = G_p(x) \cup y$ for any $x \in H^p(F, \mathbb{C})$ and $y \in H^q(M, \mathbb{C})$. The element $G_0(1) \in H^{2(n-k)}(M, \mathbb{R})$ is just the cohomology class $\text{cl}(\phi_* F)$ of the cycle $\phi_* F = \deg(F \rightarrow \phi(F))\phi(F)$. Thus,

$$\int_F (h^m)^*(\phi^* \eta^l) \wedge \phi^* \eta^{k-l} = \int_F \phi^*((f^m)^* \eta^l) \wedge \phi^* \eta^{k-l} = \int_M [(f^m)^* \eta^l] \wedge \eta^{k-l} \cup \text{cl}(\phi_* F)$$

for any Kähler form η and for any $m \geq 1$. Since ϕ is generically finite, $[\phi^* \eta] - [T] \in P^1(F)$ for a d -closed real positive $(1, 1)$ -current T by Lemma A.5. Thus, we have

$$(A-5) \quad \begin{aligned} d_l(h) &= \lim_{m \rightarrow \infty} \left(\int_M [(f^m)^* \eta^l] \wedge \eta^{k-l} \cup \text{cl}(\phi_* F) \right)^{1/m} \\ &= \lim_{m \rightarrow \infty} (\delta_l(f^m, \eta; \text{cl}(\phi_* F)))^{1/m} \leq d_l(f) \end{aligned}$$

by (2) of Lemma A.4, Lemma A.7, and (A-4). From $\phi_* \circ h_* = f_* \circ \phi_*$, we have

$$(\deg h)^{-1} G_i \circ (h^*)^{(i,i)} = (\deg f)^{-1} (f^*)^{(n-k+i, n-k+i)} \circ G_i$$

for any i , since $(\deg h)^{-1} h^*$ is the inverse of

$$H^{2i}(F, \mathbb{C}) \simeq H_{2k-2i}(F, \mathbb{C}) \xrightarrow{h_*} H_{2k-2i}(F, \mathbb{C}) \simeq H^{2i}(F, \mathbb{C})$$

and $(\deg f)^{-1} f^*$ is the inverse of

$$H^{2(n-k+i)}(M, \mathbb{C}) \simeq H_{2k-2i}(M, \mathbb{C}) \xrightarrow{f_*} H_{2k-2i}(M, \mathbb{C}) \simeq H^{2(n-k+i)}(M, \mathbb{C}).$$

We can find an eigenvector w of $(h^*)^{(l,l)}$ with the eigenvalue $d_l(h)$ from the cone $\overline{P^l(F)}$ by Theorem A.1. Thus, $f^*G_l(w) = \deg(f) \deg(h)^{-1} d_l(h) G_l(w)$. We have

$$\int_M G_l(w) \cup [\eta]^{k-l} = \int_F w \cup [\phi^* \eta]^{k-l} > 0$$

for the Kähler form η above by the projection formula for G_l and by Lemma A.4. Thus, $G_l(w) \neq 0$ and $\deg(f) \deg(h)^{-1} d_l(h)$ is an eigenvalue of $(f^*)^{(n-k+l, n-k+l)}$. Therefore, $d_{n-k+l}(f) \geq \deg(f) \deg(h)^{-1} d_l(h)$. $\square$

Now, we are ready to prove the first assertion of Theorem D.

Proof of (1) of Theorem D. We set $d = \dim Y > 0$. We have $d_1(f) \geq d_1(f|_F)$ by Proposition A.11. If $d_1(f) \leq 1$, then $\deg f = d_1(f) = d_1(f|_F) = 1$ by Fact A.10. Thus, we may assume that $d_1(f) > 1$. Let $v \in \overline{P^1(X)} \subset H^{1,1}(X, \mathbb{R})$ be an eigenvector of f^* with the eigenvalue $d_1(f)$. If $v|_F \in H^{1,1}(F, \mathbb{R})$ is not zero, then $d_1(f) \leq d_1(f|_F)$; hence $d_1(f) = d_1(f|_F)$ by Proposition A.11. Thus, it is enough to prove $v|_F \neq 0$.

There is a bimeromorphic morphism $\nu: S \rightarrow Y$ from a compact Kähler manifold S by [53]. Let $M \rightarrow X \times_Y S$ be a proper morphism from a compact Kähler manifold M , giving rise to a bimeromorphic morphism to the main component of $X \times_Y S$, i.e., the unique component dominating S . Let $\mu: M \rightarrow X$ be the induced bimeromorphic morphism and let $\varpi: M \rightarrow S$ be the induced fiber space. We may replace F with a general fiber of π , since $d_1(f|_F)$ depends only on the cohomology class $\text{cl}(F) \in H^{d,d}(X, \mathbb{R})$ (cf. (A-5)). Hence we may assume that ν is an isomorphism over a neighborhood of $\pi(F)$ and that μ is an isomorphism along $\mu^{-1}(F)$. Let $f_M = \mu^{-1} \circ f \circ \mu: M \rightarrow M$ be the meromorphic endomorphism and let $\varphi: Z \rightarrow M$ be a bimeromorphic morphism from another compact Kähler manifold Z such that $g := f_M \circ \varphi: Z \rightarrow M$ is holomorphic. Let $(f_M^*)^{(i,i)}$ be an endomorphism of $H^{i,i}(M, \mathbb{R})$ for $0 \leq i \leq n = \dim X$ defined as

$$(f_M^*)^{(i,i)}: H^{i,i}(M, \mathbb{R}) \xrightarrow{g^*} H^{i,i}(Z, \mathbb{R}) \xrightarrow{\varphi_*} H^{i,i}(M, \mathbb{R}).$$

Since $\varphi_* \circ \varphi^* = \text{id}$, we have a commutative diagram

$$\begin{array}{ccc} H^{i,i}(X, \mathbb{R}) & \xrightarrow{f^*} & H^{i,i}(X, \mathbb{R}) \\ \mu^* \downarrow & & \mu^* \downarrow \\ H^{i,i}(M, \mathbb{R}) & \xrightarrow{(f_M^*)^{(i,i)}} & H^{i,i}(M, \mathbb{R}). \end{array}$$

Thus, $\mu^* v \in \overline{P^1(M)}$ is also an eigenvector of $(f_M^*)^{(1,1)}$ with the eigenvalue $d_1(f)$. We have also $f_M^* \circ \varpi^* = \varpi^*$ from $\varpi \circ f_M = \varpi$. For any $\alpha \in H^{1,1}(X, \mathbb{R})$, $\beta \in H^{1,1}(S, \mathbb{R})$, and

$\xi \in H^{i,i}(M, \mathbb{R})$, we have the equalities

$$(A-6) \quad \begin{aligned} (f_M^*)^{(i+1,i+1)}(\xi \cup \mu^* \alpha) &= (f_M^*)^{(i,i)}(\xi) \cup \mu^* f^* \alpha, \\ (f_M^*)^{(i+1,i+1)}(\xi \cup \varpi^* \beta) &= (f_M^*)^{(i,i)} \xi \cup \varpi^* \beta \end{aligned}$$

by applying the projection formula to

$$\varphi_*(g^*(\xi \cup \mu^* \alpha)) = \varphi_*(g^* \xi \cup \varphi^* \mu^* f^* \alpha), \quad \text{and} \quad \varphi_*(g^*(\xi \cup \varpi^* \beta)) = \varphi_*(g^* \xi \cup \varphi^* \varpi^* \beta).$$

Assuming $v|_F = 0$, we shall derive a contradiction by an argument similar to [57, Section 2.1, Remark (11)]. We set $x := \mu^* \alpha$ and $y := \varpi^* \beta \in H^{1,1}(M, \mathbb{R})$ for $\alpha \in P^1(X)$ and $\beta \in P^1(S)$. Then $y^d = c \operatorname{cl}(\mu^{-1} F) \in H^{d,d}(M, \mathbb{R})$ for some $c > 0$. Thus, $\mu^* v \cup y^d = 0$. Let $1 \leq s \leq d$ be the minimum integer such that $\mu^* v \cup y^s = 0$. By [12, Corollaire 3.5], there is a constant b such that

$$(b\mu^* v + y) \cup y^{s-1} \cup \mu^*(\alpha_1 \cup \cdots \cup \alpha_{n-s-1}) = 0 \in H^{n-1,n-1}(M, \mathbb{R})$$

for any $\alpha_i \in H^{1,1}(X, \mathbb{R})$, where $n = \dim X$. Taking f_M^* , by (A-6), we have

$$(bd_1(f)\mu^* v + y) \cup y^{s-1} \cup \mu^*(f^* \alpha_1 \cup \cdots \cup f^* \alpha_{n-s-1}) = 0.$$

In particular,

$$(b\mu^* v + y) \cup y^{s-1} \cup x^{n-s} = (bd_1(f)\mu^* v + y) \cup y^{s-1} \cup x^{n-s} = 0.$$

If $b = 0$, then $y^s \cup x^{n-s} = 0$ contradicting Lemmas A.4 and A.5, since $y^s \neq 0$. Therefore, $b \neq 0$ and $\mu^* v \cup y^{s-1} \cup x^{n-s} = 0$. Then $\mu^* v \cup y^{s-1} = 0$ by Lemmas A.4 and A.5. This contradicts the minimality of s . Thus, we have completed the proof of (1) of Theorem D. $\square$

A.3. Topological entropies. Let $f: M \rightarrow M$ be a surjective endomorphism of a compact Kähler manifold M . We consider the properties of topological entropy $h_{\text{top}}(f)$ of f . Instead of giving the definition of h_{top} , we use the following:

Fact A.12 (Gromov [24], Yomdin [56] (cf. [16])). For the topological entropy $h_{\text{top}}(f)$ of f , one has

$$h_{\text{top}}(f) = \max_{1 \leq i \leq n} \log d_i(f) = \log \rho(H^*(M, \mathbb{C}), f^*).$$

As a consequence of Proposition A.11, we have:

Corollary A.13. *In the situation of Proposition A.11,*

$$h_{\text{top}}(f) \geq \log(\deg(f) \deg(h)^{-1}) + h_{\text{top}}(h) \geq h_{\text{top}}(h).$$

Notation A.14. Let $\varphi: V \rightarrow V$ be an endomorphism of a finite-dimensional $\mathbb{C}$ -vector space V . We denote by $\Lambda(V, \varphi)$ the set of eigenvalues of φ . For $\lambda \in \mathbb{C}$, we set

$$V_\lambda = V_{\lambda, \varphi} := \bigcup_{l \geq 1} \text{Ker}(\lambda \text{id}_V - \varphi)^l.$$

Remark. If $\lambda \in \Lambda(V, \varphi)$, then V_λ is a generalized eigenspace. We have the decomposition $V = \bigoplus_{\lambda \in \Lambda(V, \varphi)} V_{\lambda, \varphi}$, which is functorial in the following sense: Let $h: V_1 \rightarrow V_2$ be a $\mathbb{C}$ -linear map of finite-dimensional $\mathbb{C}$ -vector spaces. Let $\varphi_i: V_i \rightarrow V_i$ be an endomorphism for $i = 1, 2$ such that $h \circ \varphi_1 = \varphi_2 \circ h$. Then $h = \bigoplus h_\lambda$ for $h_\lambda: (V_1)_{\lambda, \varphi_1} \rightarrow (V_2)_{\lambda, \varphi_2}$.

Lemma A.15. *Let Y be a reduced compact complex analytic space and $\mu: X \rightarrow Y$ a proper surjective morphism from another reduced compact complex analytic space X such that the restriction $\mu^{-1}(U) \rightarrow U$ of μ is a smooth Kähler morphism for a dense Zariski-open subset $U \subset Y$. Let $g: Y \rightarrow Y$ and $f: X \rightarrow X$ be surjective endomorphisms such that $\mu \circ f = g \circ \mu$ and $g^{-1}(U) = U$. Let $g_B: B \rightarrow B$ and $f_A: A \rightarrow A$ be the induced endomorphisms for the complement $B = Y \setminus U$ and $A = \mu^{-1}(B)$, respectively. Then one has the following inclusion for any $p \geq 0$:*

$$\Lambda(H^p(Y, \mathbb{C}), g^*) \subset \Lambda(H^p(X, \mathbb{C}), f^*) \cup \Lambda(H^p(B, \mathbb{C}), g_B^*) \cup \Lambda(H^{p-1}(A, \mathbb{C}), f_A^*).$$

Proof. We have a commutative diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_c^i(U, \mathbb{Z}) & \longrightarrow & H^i(Y, \mathbb{Z}) & \longrightarrow & H^i(B, \mathbb{Z}) \longrightarrow \cdots \\ & & \mu^* \downarrow & & \mu^* \downarrow & & \mu|_A^* \downarrow \\ \cdots & \longrightarrow & H_c^i(\mu^{-1}(U), \mathbb{Z}) & \longrightarrow & H^i(X, \mathbb{Z}) & \longrightarrow & H^i(A, \mathbb{Z}) \longrightarrow \cdots \end{array}$$

of exact sequences, which are compatible with the actions of g^* and f^* . Since $\mu^{-1}(U) \rightarrow U$ is a smooth Kähler morphism, we have a quasi-isomorphism

$$\mathbb{R}\mu_* \mathbb{C}_X|_U \sim_{\text{qis}} \bigoplus \mathbb{R}^i \mu_* \mathbb{C}_X|_U[-i]$$

in the derived category on U , by the hard Lefschetz theorem on fibers and by [11]. In particular, the homomorphism $\mu^*: H_c^i(U, \mathbb{C}) \rightarrow H_c^i(\mu^{-1}(U), \mathbb{C})$ is injective for any i . For a complex number λ , if $H^p(Y, \mathbb{C})_{\lambda, g^*} \neq 0$ and if $H^p(B, \mathbb{C})_{\lambda, g_B^*} = H^p(X, \mathbb{C})_{\lambda, f^*} = 0$, then $H^{p-1}(A, \mathbb{C})_{\lambda, f_A^*} \neq 0$ by the commutative diagram above. Thus, the assertion follows. $\square$

Corollary A.16. *Let Y be a reduced compact complex analytic space with a surjective endomorphism $g: Y \rightarrow Y$. Then there exist a finite set $\{Z_i\}_{i=1}^N$ of closed subvarieties and a positive integer k such that $Y = \bigcup Z_i$, $(g^k)^{-1}(Z_i) = Z_i$ for any $1 \leq i \leq N$, and that the following inclusion holds for any $p \geq 0$:*

$$\Lambda(H^p(Y, \mathbb{C}), (g^k)^*) \subset \bigcup_{i=1}^N \bigcup_{q=0}^p \Lambda(H^q(Z_i, \mathbb{C}), (g^k|_{Z_i})^*).$$

Proof. Let $\{Y_i\}$ be the set of the irreducible components of Y and let $\tau: X := \bigsqcup Y_i \rightarrow Y$ be the natural morphism. Let $B \subset Y$ be the minimum closed subset such that $X \setminus \tau^{-1}B \rightarrow Y \setminus B$ is an isomorphism. By replacing g with a power g^k , we may assume that $g^{-1}Y_i = Y_i$ for any i . Then, an endomorphism $f: X \rightarrow X$ with $\tau \circ f = g \circ \tau$ is induced, where $g^{-1}(B) = B$ by the minimality of B and $f^{-1}(\tau^{-1}(B)) = \tau^{-1}(B)$. Applying Lemma A.15 to $X \rightarrow Y$, f , g , B , and $\tau^{-1}(B)$, we have the following inclusion for any p :

$$\Lambda(H^p(Y, \mathbb{C}), g^*) \subset \Lambda(H^p(B, \mathbb{C}), g^*) \cup \bigcup \left(\Lambda(H^p(Y_i, \mathbb{C}), g^*) \cup \Lambda(H^{p-1}(Y_i \cap B, \mathbb{C}), g^*) \right).$$

Applying the same argument to B and $Y_i \cap B$, and continuing, we have the assertion. $\square$

Proposition A.17. *Let Y be a reduced compact complex analytic space and let $\phi: M \rightarrow Y$ be a proper surjective morphism from a finite disjoint union M of compact Kähler manifolds M_α such that $\phi(M_\alpha)$ is an irreducible component of Y . Let $g: Y \rightarrow Y$ and $f: M \rightarrow M$ be étale surjective endomorphisms such that $\phi \circ f = g \circ \phi$. If $f^{-1}(M_\alpha) = M_\alpha$ for any α , then, for the induced endomorphisms $f|_{M_\alpha}: M_\alpha \rightarrow M_\alpha$, one has*

$$\rho(H^*(Y, \mathbb{C}), g^*) \leq \max_\alpha \rho(H^*(M_\alpha, \mathbb{C}), (f|_{M_\alpha})^*) = \max_{\alpha, l} d_l(f|_{M_\alpha}).$$

Proof. We shall prove by induction on $\dim M = \max\{\dim M_\alpha\}$. If $\dim M = 0$, then the assertion holds by a trivial reason. Assume that the assertion holds when the dimension is less than $\dim M$. If the inequality of the spectral radius holds for a power g^l , then it holds also for g ; in fact, $\rho(H^p(Y, \mathbb{C}), (g^l)^*) = \rho(H^p(Y, \mathbb{C}), (g^*)^l)$. Thus, we may replace g with any power g^l . Let $U \subset Y$ be the maximal Zariski open subset such that $\phi^{-1}(U) \rightarrow U$ is smooth. For the complement $B = Y \setminus U$ and $A = \phi^{-1}(B)$, we have $g^{-1}(B) = B$ and $f^{-1}(A) = A$ by the maximality of U and by the étaleness of g and f . Thus, we can apply Lemma A.15, and obtain

$$\rho(H^*(Y, \mathbb{C}), g^*) \leq \max\{\rho(H^*(M, \mathbb{C}), f^*), \rho(H^*(B, \mathbb{C}), (f|_B)^*), \rho(H^*(A, \mathbb{C}), (f|_A)^*)\}.$$

Let $A = \bigcup A_\beta$ be the irreducible decomposition. Replacing g with its power, we may assume that $f^{-1}(A_\beta) = A_\beta$ for any β . Let $Z_\beta \rightarrow A_\beta$ be an equivariant resolution of singularities of A_β with respect to the étale endomorphism $f|_{A_\beta}$. Let Z be the disjoint union $\bigsqcup Z_\beta$ and let $h: Z \rightarrow Z$ be the induced étale endomorphism, i.e., $\psi \circ h = f|_A \circ \psi$ for the induced morphism $\psi: Z \rightarrow A$. Then $(Z \rightarrow A \rightarrow B, h, g|_B)$ and $(Z \rightarrow A, h, f|_A)$ satisfy the same condition of Proposition A.17 as $(M \rightarrow Y, f, g)$. By induction, we have

$$\max\{\rho(H^*(B, \mathbb{C}), (g|_B)^*), \rho(H^*(A, \mathbb{C}), (f|_A)^*)\} \leq \max_\beta \rho(H^*(Z_\beta, \mathbb{C}), (h|_{Z_\beta})^*)$$

after replacing g with a power g^l . On the other hand,

$$\rho(H^*(Z_\beta, \mathbb{C}), (h|_{Z_\beta})^*) = \exp(h_{\text{top}}(h|_{Z_\beta})) \leq \exp(h_{\text{top}}(f|_{M_\alpha})) = \rho(H^*(M_\alpha, \mathbb{C}), (f|_{M_\alpha})^*)$$

for $A_\beta \subset M_\alpha$, by Proposition A.11. Hence,

$$\rho(H^*(Y, \mathbb{C}), g^*) \leq \max_\alpha \rho(H^*(M_\alpha, \mathbb{C}), (f|_{M_\alpha})^*). \quad \square$$

Lemma A.18. *Let $\pi: X \rightarrow Y$ be a smooth fiber space of complex manifolds and $f: X \rightarrow X$ a surjective endomorphism such that $\pi \circ f = \pi$. Let F be a fiber of π . Then, for any $p \geq 0$, one has an inclusion*

$$\Lambda(H_c^p(X, \mathbb{C}), f^*) \subset \bigcup_{q=0}^p \Lambda(H^q(F, \mathbb{C}), (f|_F)^*).$$

Proof. We have an endomorphism f^* of the complex $R\pi_*\mathbb{C}_X$ in the derived category of sheaves of abelian groups on Y such that the induced endomorphism of $R\Gamma_c(Y, R\pi_*\mathbb{C}_X)$ coincides with that of $R\Gamma_c(X, \mathbb{C})$. In particular, the Leray spectral sequence

$$E_2^{p,q} = H_c^p(Y, R^q\pi_*\mathbb{C}_X) \Rightarrow E^{p+q} = H_c^{p+q}(X, \mathbb{C})$$

admits the endomorphisms $f^*: E_r^{p,q} \rightarrow E_r^{p,q}$ compatible with f^* on E^{p+q} . For a complex number λ , let $(R^i\pi_*\mathbb{C}_X)_\lambda$ be the union of the kernels of

$$(f^* - \lambda \text{id})^l: R^i\pi_*\mathbb{C}_X \rightarrow R^i\pi_*\mathbb{C}_X$$

for all $l > 0$ (cf. Notation A.14). Then,

$$(E_2^{p,q})_\lambda \simeq H_c^p(Y, (R^q\pi_*\mathbb{C}_X)_\lambda)$$

for any p, q , and λ . Since $E_2^{p,q} \Rightarrow E^{p+q}$ is decomposed into $(E_2^{p,q})_\lambda \Rightarrow (E^{p+q})_\lambda$, we have

$$\Lambda(H_c^p(X, \mathbb{C}), f^*) \subset \bigcup_{q=0}^p \{\lambda \in \mathbb{C} \mid (R^q\pi_*\mathbb{C}_X)_\lambda \neq 0\} = \bigcup_{q=0}^p \Lambda(H^q(F, \mathbb{C}), (f|_F)^*). \quad \square$$

We are ready to prove the second assertion of Theorem D.

Proof of (2) of Theorem D. We may replace f with a power f^k since $h_{\text{top}}(f^k) = kh_{\text{top}}(f)$ and $h_{\text{top}}(f^k|_F) = kh_{\text{top}}(f|_F)$.

Step 1. The first reduction: Let $\nu: Y' \rightarrow Y$ be a bimeromorphic morphism from a compact Kähler manifold Y' . Then f induces an étale endomorphism $f \times_Y Y': X \times_Y Y' \rightarrow X \times_Y Y'$. Let X' be the main component of $X \times_Y Y'$. Thus, there exists an equivariant resolution of singularities $\widetilde{X} \rightarrow X'$ with respect to the étale endomorphism. We may assume $\widetilde{X}$ to be Kähler since $\widetilde{X} \rightarrow X'$ can be chosen as a projective morphism. Let $\tilde{f}$ be the induced étale endomorphism of $\widetilde{X}$. Then $d_l(\tilde{f}) = d_l(f)$ by Lemma A.8. Thus, we may assume that Y is also a compact Kähler manifold and $\pi: X \rightarrow Y$ is smooth over the complement $Y \setminus D$ for a divisor $D \subset Y$.

Step 2: A setting of the proof. Let $\sigma: S \rightarrow Y$ be a flattening of π , i.e., the main component of $X \times_Y S$ is flat over S (cf. [29]). Here, we may assume that $\sigma^{-1}D$ is also a normal crossing divisor. Let $M \rightarrow X \times_Y S$ be a bimeromorphic morphism from a compact Kähler manifold M which is given as an equivariant resolution of singularities of the main

component with respect to the induced étale endomorphism $f \times_Y \text{id}_S: X \times_Y S \rightarrow X \times_Y S$. Let $\mu: M \rightarrow X$ be the induced bimeromorphic morphism, $\varpi: M \rightarrow S$ the induced fiber space, and $g: M \rightarrow M$ the induced étale endomorphism, i.e., $\mu \circ g = f \circ \mu$ and $\varpi \circ g = \varpi$. We set $E = \pi^*(D)_{\text{red}}$. For the prime decomposition $E = \sum_{j=1}^l E_j$, there is a positive integer k such that $(f^k)^{-1}E_j = E_j$. Here, we may assume $k = 1$ by replacing f with f^k ; hence $f^{-1}(E_j) = E_j$. Let G_j be the proper transform of E_j in M for any j . Then $g^{-1}(G_j) = G_j$. By a suitable choice of an equivariant embedded resolution of singularities, we may assume that all G_j are nonsingular. Then $\varpi(G_j)$ is a prime component of $\sigma^{-1}D$ for any j by the flattening. Let $h_j: G_j \rightarrow G_j$ be the étale endomorphism $g|_{G_j}$. In order to prove (2) of Theorem D, it is enough to show the following two inequalities:

$$(A-7) \quad h_{\text{top}}(f) \leq \max\{h_{\text{top}}(f|_F), h_{\text{top}}(h_j) \mid 1 \leq j \leq l\}.$$

$$(A-8) \quad h_{\text{top}}(f|_F) \geq h_{\text{top}}(h_j) \quad \text{for any } 1 \leq j \leq l.$$

Indeed, we have already the other inequality $h_{\text{top}}(f) \geq h_{\text{top}}(f|_F)$ in Corollary A.13. The inequality (A-7) is shown in *Step 3* below, while the other inequality (A-8) is shown in *Step 4* by using induction on the dimension of X .

Step 3. Proof of (A-7): The natural long exact sequence

$$\cdots \rightarrow H_c^p(X \setminus E, \mathbb{C}) \rightarrow H^p(X, \mathbb{C}) \rightarrow H^p(E, \mathbb{C}) \rightarrow \cdots$$

admits an endomorphism f^* induced from f . Thus,

$$\begin{aligned} \Lambda(H^*(X, \mathbb{C}), f^*) &\subset \Lambda(H_c^*(X \setminus E, \mathbb{C}), (f|_{X \setminus E})^*) \cup \Lambda(H^*(E, \mathbb{C}), (f|_E)^*) \\ &\subset \Lambda(H^*(F, \mathbb{C}), (f|_F)^*) \cup \bigcup_j \Lambda(H^*(G_j, \mathbb{C}), h_j^*) \end{aligned}$$

by Proposition A.17 applied to $\sqcup G_j \rightarrow E$ and Lemma A.18 applied to $X \setminus E \rightarrow Y \setminus D$. Therefore, (A-7) follows.

Step 4. Proof of (A-8): By induction, we may assume that (2) of Theorem D holds for lower dimensional ambient spaces. Let P be a general point of $\varpi(G_j)$. We may assume that P is a nonsingular point of $\sigma^{-1}D$, ϖ is flat over an open neighborhood of P , and $G_j \rightarrow \varpi(G_j)$ is smooth over an open neighborhood of P . Let $G_j \rightarrow S_j$ be the fiber space obtained as the Stein factorization of $G_j \rightarrow \varpi(G_j)$. Let Γ be the fiber of $G_j \rightarrow S_j$ over a point of S_j lying over P . By induction, we may apply (2) of Theorem D to the fiber space $G_j \rightarrow S_j$ and the étale endomorphism $g|_{G_j}$. Thus, $h_{\text{top}}(h_j) = h_{\text{top}}(g|_{\Gamma})$. Let $\text{cl}(\Gamma)$ and $\text{cl}(\varpi^{-1}(P))$ be the cohomology classes in $H^{d,d}(M, \mathbb{R})$ corresponding to the subspaces Γ and $\varpi^{-1}(P)$, respectively, where $d = \dim S = \dim Y$. Then $\text{cl}(\varpi^{-1}(P)) - \text{cl}(\Gamma)$ is represented by a positive current. The class $\text{cl}(\varpi^{-1}(P)) \in H^{d,d}(M, \mathbb{R})$ equals the class $\text{cl}(F')$ corresponding to a smooth fiber F' of $\varpi: M \rightarrow S$, since ϖ is flat along $\varpi^{-1}(P)$.

Thus,

$$\begin{aligned}\delta_l((g|_\Gamma)^m, \eta|_\Gamma) &= \delta_l(g^m, \eta; \text{cl}(\Gamma)) \leq \delta_l(g^m, \eta; \text{cl}(\varpi^{-1}(P))) \\ &= \delta_l(g^m, \eta; \text{cl}(F')) = \delta_l((g|_{F'})^m, \eta|_{F'})\end{aligned}$$

for a Kähler form η , any $0 \leq l \leq n - d$, and any $m \geq 1$. In particular, $h_{\text{top}}(g|_\Gamma) \leq h_{\text{top}}(g|_{F'})$. We may assume that $\mu: M \rightarrow X$ is an isomorphism along F' and $\sigma: S \rightarrow Y$ is an isomorphism at the point $\varpi(F')$. Therefore, $h_{\text{top}}(h_j) = h_{\text{top}}(g|_\Gamma) \leq h_{\text{top}}(g|_{F'}) = h_{\text{top}}(f|_F)$. Thus, we have completed the proof of (2) of Theorem D. $\square$

REFERENCES

- [1] E. Amerik and F. Campana, Fibrations méromorphes sur certaines variétés de classe canonique triviale, preprint 2005 (math.AG/0510299).
- [2] B. Angéniol, *Familles de Cycle Algébriques – Schéma de Chow*, Lecture Notes in Math., **896**, Springer, 1981.
- [3] D. Barlet, Espace analytique réduit des cycles analytiques complexes compacts d'un espace analytique complexe de dimension finie, *Fonctions de plusieurs variables complexes, II (Sém. François Norguet, 1974–1975)*, pp. 1–158, Lecture Notes in Math., **482**, Springer, 1975.
- [4] A. Beauville, Some remarks on Kähler manifolds with $c_1 = 0$, *Classification of Algebraic and Analytic Manifolds* (Katata, 1982, ed. K. Ueno), Progr. Math., **39** Birkhäuser 1983, pp. 1–26.
- [5] C. Birkar, P. Cascini, C. D. Hacon and J. McKernan, Existence of minimal models for varieties of log general type, preprint 2006 (math.AG/0610203).
- [6] E. Bierstone and P. D. Milman, Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant, *Invent. Math.*, **128** (1997), 207–302.
- [7] G. Birkhoff, Linear transformations with invariant cones, *Amer. Math. Monthly*, **74** (1967), 274–276.
- [8] F. Bogomolov, On the decomposition of Kähler manifolds with trivial canonical class, *Math. USSR Sbornik*, **22** (1974), 580–583.
- [9] F. Campana, On twister spaces of the class $\mathcal{C}$, *J. Diff. Geom.* **33** (1991), 541–549.
- [10] F. Campana, Connexité rationnelle des variétés de Fano, *Ann. Sci. École Norm. Sup. (4)* **25** (1992), 539–545.
- [11] P. Deligne, Théorème de Lefschetz et critères de dégénérescence de suites spectrales, *Publ. Math. Inst. Hautes Études Sci.*, **35** (1968), 107–126.
- [12] T.-C. Dinh and N. Sibony, Groupes commutatifs d'automorphismes d'une variété kählérienne compacte, *Duke Math. J.* **123** (2004), 311–328.
- [13] T.-C. Dinh and N. Sibony, Green currents for holomorphic automorphisms of compact Kähler manifolds, *J. Amer. Math. Soc.* **18** (2005), 291–312.
- [14] S. Encinas and H. Hauser, Strong resolution of singularities in characteristic zero, *Comment. Math. Helv.*, **77** (2002), 821–845.
- [15] S. Encinas and O. E. Villamayor, A course on constructive desingularization and equivariance, *Resolution of singularities (Obergrugl, 1997)*, Progr. Math. **181**, Birkhäuser, 2000, pp. 147–227.
- [16] S. Friedland, Entropy of polynomial and rational maps, *Ann. of Math.* **133** (1991), 359–368.

- [17] A. Fujiki, On the blowing down of analytic spaces, Publ. Res. Inst. Math. Sci. Kyoto Univ., **10** (1975), 473–507.
- [18] A. Fujiki, Closedness of the Douady spaces of compact Kähler spaces, Publ. Res. Inst. Math. Sci. Kyoto Univ., **14** (1978), 1–52.
- [19] A. Fujiki, On automorphism groups of compact Kähler manifolds, Invent. Math., **44** (1978), 225–258.
- [20] Y. Fujimoto, Endomorphisms of smooth projective 3-folds with nonnegative Kodaira dimension, Publ. Res. Inst. Math. Sci. Kyoto Univ., **38** (2002), 33–92.
- [21] Y. Fujimoto and N. Nakayama, Endomorphisms of smooth projective 3-folds with nonnegative Kodaira dimension, II, J. Math. Kyoto Univ., **47** (2007), 79–114.
- [22] T. Graber, J. Harris and J. Starr, Families of rationally connected varieties, J. Amer. Math. Soc. **16** (2003), 57–67.
- [23] P. Griffiths, Periods of integrals on algebraic manifolds, III, Publ. Math. I.H.E.S. **38** (1970), 125–180.
- [24] M. Gromov, On the entropy of holomorphic maps, Enseign. Math. **49** (2003), 217–235.
- [25] V. Guedj, Ergodic properties of rational mappings with large topological degree, Ann. of Math. **161** (2005), 1589–1607.
- [26] V. Guedj, Propriétés ergodiques des applications rationnelles, preprint 2006 (math.CV/0611302v1).
- [27] M. Hanamura, On the birational automorphism groups of algebraic varieties, Compos. Math. **63** (1987), 123–142.
- [28] M. Hanamura, Structure of birational automorphism groups, I: non-uniruled varieties, Invent. Math., **93** (1988), 383–403.
- [29] H. Hironaka, Flattening theorem in complex-analytic geometry, Amer. J. Math., **97** (1975), 503–547.
- [30] S. Iitaka, On logarithmic Kodaira dimension of algebraic varieties, *Complex analysis and algebraic geometry* (eds. W. L. Baily, Jr. and T. Shioda), pp. 175–189, Iwanami Shoten and Cambridge Univ. Press, 1977.
- [31] S. Iitaka, *Algebraic Geometry – an Introduction to Birational Geometry of Algebraic Varieties*, Grad. Texts in Math., **76**, Springer 1981.
- [32] Y. Kawamata, Characterization of abelian varieties, Compos. Math. **43** (1981), 253–276.
- [33] Y. Kawamata, Minimal models and the Kodaira dimension of algebraic fiber spaces, J. Reine Angew. Math. **363** (1985), 1–46.
- [34] Y. Kawamata, Abundance theorem for minimal threefolds, Invent. Math., **108** (1992), 229–246.
- [35] Y. Kawamata, K. Matsuda and K. Matsuki, Introduction to the minimal model problem, *Algebraic geometry, Sendai, 1985* (T. Oda ed.), Adv. Stud. Pure Math., **10**, Kinokuniya and North-Holland, 1987, pp. 283–360.
- [36] J. Kollár, Higher direct images of dualizing sheaves, II, Ann. of Math., **124** (1986), 171–202.
- [37] J. Kollár, Shararevich maps and plurigenera of algebraic varieties, Invent. Math., **113** (1993), 177–215.
- [38] J. Kollár, *Rational curves on algebraic varieties*, Ergeb. Math. Grenzgeb. (3) **32**, Springer, 1996.
- [39] J. Kollár, *Lectures on resolution of singularities*, Ann. of Math. Studies, **166**, Princeton University Press, 2007.
- [40] J. Kollár, Y. Miyaoka and S. Mori, Rational connected varieties, J. Alg. Geom. **1** (1992), 429–448.
- [41] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., **134**, Cambridge Univ. Press, 1998.

- [42] D. I. Lieberman, Compactness of the Chow scheme: applications to automorphisms and deformations of Kähler manifolds, *Fonctions de plusieurs variables complexes, III (Sém. François Norguet, 1975–1977)*, pp. 140–186, Lecture Notes in Math., **670**, Springer, 1978.
- [43] Y. Miyaoka, On the Kodaira dimension of minimal threefolds, *Math. Ann.*, **281** (1988), 325–332.
- [44] Y. Miyaoka, Abundance conjecture for 3-folds: case $\nu = 1$, *Compos. Math.*, **68** (1988), 203–220.
- [45] Y. Miyaoka and S. Mori, A numerical criterion for uniruledness, *Ann. of Math.*, **124** (1986), 65–69.
- [46] S. Mori, Classification of higher-dimensional varieties, *Algebraic geometry, Bowdoin, 1985*, pp. 269–331, *Proc. Sympos. Pure Math.*, **46**, Amer. Math. Soc., 1987.
- [47] I. Nakamura and K. Ueno, An addition formula for Kodaira dimension of analytic fibre bundles whose fibre are Moisèzon manifolds, *J. Math. Soc. Japan* **25** (1973), 363–371.
- [48] N. Nakayama, Compact Kähler manifolds whose universal covering spaces are biholomorphic to $\mathbb{C}^n$, preprint RIMS-**1230**, Res. Inst. Math. Sci. Kyoto Univ. 1999; currently in the midst of revision.
- [49] N. Nakayama, Ruled surfaces with non-trivial surjective endomorphisms, *Kyushu J. Math.*, **56** (2002), 433–446.
- [50] Y. Namikawa and J. Steenbrink, Global smoothing of Calabi-Yau threefolds, *Invent. Math.*, **122** (1995), 403–419.
- [51] S. Takayama, Local simple connectedness of resolutions of log-terminal singularities, *Internat. J. Math.* **14** (2003), 825–836.
- [52] K. Ueno, *Classification theory of algebraic varieties and compact complex spaces*, Lecture Notes in Mathematics, **439**, Springer, 1975.
- [53] J. Varouchas, Stabilité de la classe des variétés kähleriennes pour les certains morphismes propres, *Invent. Math.*, **77** (1984), 117–128.
- [54] E. Viehweg and K. Zuo, On the isotriviality of families of projective manifolds over curves, *J. Alg. Geom.* **10** (2001), 781–799.
- [55] J. Włodarczyk, Simple Hironaka resolution in characteristic zero, *J. Amer. Math. Soc.*, **18** (2005), 779–822.
- [56] Y. Yomdin, Volume growth and entropy, *Israel J. Math.* **57** (1987), 285–300.
- [57] D.-Q. Zhang, Dynamics of automorphisms on projective complex manifolds, *J. Diff. Geom.* (to appear); also: arXiv:0810.4675v1.

(Noboru Nakayama) RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES
 KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN
E-mail address: nakayama@kurims.kyoto-u.ac.jp

(De-Qi Zhang) DEPARTMENT OF MATHEMATICS
 NATIONAL UNIVERSITY OF SINGAPORE, 2 SCIENCE DRIVE 2, SINGAPORE 117543
E-mail address: matzdzq@nus.edu.sg